

Compilation of Giant Electric Dipole Resonances Built on Excited States

A. SCHILLER *

National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, MI 48824
and

M. THOENNESSEN

National Superconducting Cyclotron Laboratory, and
Department of Physics & Astronomy, Michigan State University, East Lansing, MI 48824

Giant Electric Dipole Resonance (GDR) parameters for γ decay to excited states with finite spin and temperature are compiled. Over 100 original works have been reviewed and from some 70 of which more than 300 parameter sets of hot GDR parameters for different isotopes, excitation energies, and spin regions have been extracted. All parameter sets have been brought onto a common footing by calculating the equivalent Lorentzian parameters. The current compilation is complementary to an earlier compilation by Samuel S. Dietrich and Barry L. Berman (At. Data Nucl. Data Tables **38**, 199-338, 1988) on ground-state photo-neutron and photo-absorption cross sections and their Lorentzian parameters. A comparison of the two may help shed light on the evolution of GDR parameters with temperature and spin. The present compilation is current as of January 2006.

* Corresponding author: NSCL/MSU, 1 Cyclotron, East Lansing, MI 48824. Phone: (517) 324-8142, Fax: (517) 353-5967
Email address: schiller@nscl.msu.edu (A. Schiller).

CONTENTS

1	INTRODUCTION	2
1.1	Theoretical considerations	2
1.2	Experimental Techniques	3
1.3	Statistical-model calculations and comparison to experimental data	5
1.4	Data treatment	6
1.5	Policy	7
1.6	Note on References	8
	EXPLANATION OF TABLE	13
	TABLE	
I.	Hot GDR parameters	15
	REFERENCES FOR TABLE	28

1. INTRODUCTION

Over the last decades, numerous data on hot Giant Electric Dipole Resonance (GDR) parameters have been accumulated. Since the field seems to have matured, it might be useful at this point to gather all present data in one comprehensive compilation in a uniform format. The Introduction is organized in the following way: first, we give some theoretical motivation why GDR parameters might be dependent on temperature, hence what question the present compilation tries to address. Then, in Subsection 1.2, we give an overview over typical experimental techniques used for measuring hot GDR parameters; in Subsection 1.3 we describe different ways of how hot GDR parameters are extracted from such experiments by means of statistical-model calculations. Although we have not attempted to extract any GDR parameters from such calculations ourselves, we believe it is helpful for the reader to get a general impression of the experimental and data-analysis part of the compiled works. In Subsection 1.4 we explain how different sets of GDR parameterizations from the original articles were brought onto a common footing. Finally, we end the Introduction with a statement regarding our policy and a note on references.

1.1. Theoretical considerations

A first good understanding of statistical γ emission was gained from the works of Brink [1] and Axel [2] who realized that average electric-dipole ($E1$) transition strengths in different energy regimes can be described in a unified fashion by assuming that the GDR can be built on any excited state, and that the GDR properties do not depend on the temperature of the excited state in question.

This so-called Brink-Axel hypothesis has been refined in the past to allow for temperature- and spin-dependent widths. A model to motivate such a modification takes into account shape fluctuations of the nucleus. Since the ground-state GDR splits into two components for a nucleus with static deformation, and the splitting depends on the degree of deformation [3], it is reasonable to assume that at finite temperatures, when the nucleus can explore a large volume in deformation space, the GDR response will be an average over different deformations and hence, different splittings. The result within this adiabatic damping model will be a more diffuse and certainly wider GDR than the ground-state GDR.

Quantitatively, assuming a Fermi-gas level density, one can write the nuclear entropy in the micro-canonical ensemble as $S = 2\sqrt{a(E - V)}$, where V is some potential energy proportional to the square of the deformation $V = k\beta^2$. The entropy is trivially maximized for $\beta = 0$; expanding S for small $V \ll E$ yields $S \approx 2\sqrt{aE} (1 - \frac{V}{2E})$. With $T = \sqrt{E/a}$, the probability distribution to find a nucleus with energy E and deformation β becomes $P \approx \exp(2\sqrt{aE}) \exp(-k\beta^2/T)$ where the second factor represents a Gaussian distribution of deformations around $\beta = 0$ and a width of $\sqrt{T/2k}$. Assuming that the splitting of the GDR into two components is roughly proportional to the nuclear deformation, the shape-fluctuation model predicts an increase in width of the GDR roughly proportional to $\sqrt{T}$. Moreover, there is also a potential spin dependence of the GDR width which stems from the possibility of spin-induced deformation. Finally, orientation fluctuations of the nucleus and nuclear-structure effects such as pairing can influence the temperature dependence of the GDR in different energy regimes. Several groups have calculated temperature-dependent GDR widths along these lines [4; 5; 6] and a simple scaling law has emerged [4].

Investigations of the low-energy tail of the GDR have also yielded indications for a temperature-dependent GDR width. It was, e.g., noted by Popov [7] in $(n, \gamma\alpha)$ experiments on Sm nuclei that the γ strength function tends to approach a finite value for $E_\gamma \rightarrow 0$ for γ transitions in the quasi-continuum (below the neutron separation energy). This experimental observation led Kadenskii, Markushev, and Furman (KMF) to propose a γ strength-function model for spherical nuclei with a temperature-dependent width [8] based on the effect of in-medium nucleon-nucleon collisions. The proposed temperature dependence was derived within Migdal's theory of Fermi liquids and has the form $\Gamma(E_\gamma, T) \propto (E_\gamma^2 + 4\pi^2 T^2)$. The model was later improved by Sirotkin [9] who included the Pauli exclusion principle, and it was extended to deformed nuclei within the framework of the generalized Lorentzian model by Kopecky and Uhl [10] and by inclusion of a coupling term between the $E1$ operator and the quadrupole deformation according to Mughabghab and Dunford [11]. The KMF model (taken at constant temperatures) and its extensions have been successfully applied to improve γ and isomeric production cross sections [12] and they have been used for direct fits of measured low-energy γ strength functions [13]. The connection of collisional-damping models with hot GDR parameters has been made in [14].

1.2. Experimental Techniques

Unlike the measurement of the GDR by ground-state photo-absorption cross-section measurements [15], measurements of hot GDR parameters can be performed in many different ways. One of the

simplest ways is by fusion-evaporation reactions where only γ rays are detected [16]. Such measurements are the most inclusive reactions, since the high-energy γ yield which competes with particle and especially neutron evaporation is representative for a range of different product nuclei, excitation energies, and spins. Moreover, it is not necessarily guaranteed that all detected γ rays stem from fusion-evaporation reactions. Other reactions such as inelastic or deep inelastic scattering can compete and yield γ rays from target or projectile-like fragments.

To improve the sensitivity of such experiments to the fusion-evaporation reaction channel, typical gates such as, e.g., γ -multiplicity filters [17], detection of heavy evaporation residues [18], and detection of evaporated, light charged particles such as protons or α particles [19] can be performed. The resulting γ -ray spectra are more exclusive, not only in terms of the product nuclei from which high-energy γ rays are emitted, but also in terms of the spin and the excitation-energy range investigated. For example, a gate on different γ folds translates rather directly into certain spin regions of the investigated product nucleus [17]. A gate on evaporated light charged particles will not only reduce the average charge and mass of the product nucleus, but it will also reduce its average excitation energy, since the evaporated particles will carry away some part of the initial excitation energy of the compound nucleus [20]; hence applying such a gate will test the GDR at somewhat lower temperatures than the fully inclusive experiment. In the same way, gating on the γ sum energy [21] or on specific product nuclei by means of detecting in coincidence discrete, known low-energy γ transitions [22] will also influence the average spin and excitation-energy region from which the high-energy γ rays are emitted, since one effectively biases the competition between high-energy γ decay and neutron evaporation in one or the other direction. Other, more rarely used gating conditions are, e.g., the isomeric γ decay by discrete transitions [23] or the α decay of a product nucleus [24], both of which have similar implications for the average spin and excitation-energy range from which prompt high-energy γ rays are observed.

For heavy nuclei, an added difficulty is the possibility of fission of the compound nucleus. Typically, for low spins, the production of an evaporation residue dominates while for high spins fission will become the dominant exit channel [25]. Hence, by gating on evaporation residues or fission fragments, one effectively selects a spin region from which high-energy γ emission is observed [26]. In the case of the fission exit channel, one also observes high-energy γ emission from the fission fragments themselves [27], though typically at significantly higher γ energies owing to the much lower mass of the fission fragments. Another complication is the fact that high-energy γ emission can occur from the compound nucleus (which is desired), or during the saddle-to-scission motion after the nucleus has passed the fission barrier [28]. Also, for the heaviest nuclei investigated, it is not clear whether a true compound nucleus forms which is confined by some fission barrier or whether one observes direct fission or just the formation of a mononucleus. Where we suspect the latter as in, e.g., [29], the extracted data do not enter the present compilation.

In some cases, one employs inelastic (see, e.g., [30]) or deep inelastic scattering [31; 32; 33; 34; 35] to excite the target or projectile nucleus. By measuring the kinetic energy of at least one of the products, one can reconstruct the reaction kinematics event by event and it is possible to obtain initial excitation-energy indexed coincident high-energy γ spectra with only one beam energy. Otherwise, the excited nucleus can be treated in the same way as a compound nucleus which is formed in a fusion-evaporation reaction.

1.3. Statistical-model calculations and comparison to experimental data

High-energy γ spectra are typically analyzed using a statistical-model calculation. In the first step, total fusion cross sections and maximum (I_0) or average ($\langle I_i \rangle$) angular momenta are determined. Typically, total fusion cross sections can be verified by experiment; maximum angular momenta are calculated by the theory of either Winther [36] or Swiatecki [37]. Average angular momenta can then be determined by $\langle I_i \rangle = \frac{2}{3}I_0$. In the next step, the decay of the highly excited compound nucleus is modeled. In many cases, this simply involves a Hauser-Feshbach-type theory [38] into which particle and γ transmission coefficients (sometimes including higher-than- $E1$ multipolarities) as well as nuclear level densities enter. Typically, particle transmission coefficients are not discussed in great detail in the compiled works. The level-density models are either the Pühlhofer model [39] (the default in the statistical-model code CASCADE [39]) or the Reisdorf model [40]. The Pühlhofer model relies on the local Dilg *et al.* parameterization [41] for excitation energies up to and slightly above the nucleon separation energy, while it interpolates then to a regime where the level-density parameter a becomes proportional to the nuclear mass number A . The Reisdorf approach builds on the generalized superfluid model by Ignatyuk *et al.* [42], but it uses a global parameterization for the asymptotic level density parameter a . In one case [30], the level-density model by Fineman *et al.* [43] is used in the data analysis.

Statistical-model calculations are often adapted to different experimental situations. For light compound nuclei near the $N = Z$ line, an isospin-dependent formalism is often used [16]. Also, the Wigner energy [44] is sometimes included in the level-density parameterization [16]. For large excitation energies, pre-equilibrium emission due to direct and semi-direct reaction mechanisms are often taken into account (see, e.g., [45]). Especially the PEQAG2 code [46] has been developed for this purpose. In the case of fissile compound nuclei, the fission channel and the decay of excited fission fragments are modeled as well (see, e.g., [25]). When gating conditions were applied in the experiment, they are usually reflected in the statistical-model calculation as well, which often implies the need to use a Monte-Carlo simulation tool such as in [19]. In some cases, also asymmetries a_2 of γ emission are calculated (see, e.g., [21]), however, experimental asymmetries do not enter into the present work (with the exception of their influence on the sign of the quoted deformation as in, e.g., [21]).

Typically, calculated high-energy γ spectra are compared to their experimental counterparts, and the γ transmission coefficients (parameterized by one- or two-component Lorentzians multiplied by a factor $2\pi E_\gamma^3$) are varied until the best fit is obtained. If absolute values in the high-energy region are compared, the fit is often normalized to the data in an energy region of 3–7 MeV (see, e.g., [47]), far below the peak of the GDR. Statistical uncertainties are usually determined in the normal fashion by varying GDR parameters until the quality of the fit deteriorates. Systematic uncertainties are more difficult to estimate. A good way is, e.g., to perform several fits to the experimental data with different level-density parameterizations as in [48] or differing sets of other input parameters into the statistical-model calculation (say, e.g., those which describe the dynamic of the fission process such as the nuclear viscosity which governs the timescale of the saddle-to-scission motion as in [25]). The range of resulting GDR parameters might give a good indication of the size of the systematic error.

Other sources of systematic errors concern the experimental conditions. Some of the most important problems there involve inefficient neutron- γ discrimination (often done by time-of-flight techniques as

in [30], or by simply considering γ rays only at backward angles as in [49]), contamination of high-energy γ spectra by cosmic rays (which can be greatly reduced by coincidence measurements as in [50]), target impurities, pile-up (see, e.g., [51] for a thorough investigation of these two effects), and add-back issues. Add back is a technique often used for an array of small detectors where for high-energy γ rays one observes a significant amount of (i) Compton scattering from one detector into a neighboring one, and (ii) pair production with subsequent annihilation γ rays being detected in neighboring detectors. The add-back technique remedies this situation by adding the deposited energies in neighboring detectors and (rightly) consider such events as stemming from one single γ ray (see, e.g., [21]). Pile up is a problem especially for large detectors, where two or more coincident γ rays hit the same detector and their energies add up and are falsely registered as one high-energy γ ray (see, e.g., the discussion in [51]). If one applies the add-back technique, however, pile up can also occur when two coincident γ rays hit two neighboring detectors. Obviously, for any given detector array, there is an optimal balance between the benefit of applying the add-back technique and the possible distortions of the high-energy γ spectrum due to pile up. Where we found that this balance was not met [52], we have rejected the data for the present compilation.

Some other physical background involves nuclear bremsstrahlung which is emitted in the first moments of the fusion process where individual nucleons of the projectile are greatly de-accelerated in the proximity of target nucleons and emit high-energy γ rays (see, e.g., the discussion in [51]). These γ rays can either be modeled (and hence subtracted from the high-energy γ spectrum which is to be fitted by the statistical-model calculation as in [30]), or they are simply considered as a source of systematic error as in [51]. Another possible source of high-energy γ rays is the pre-equilibrium γ emission during the formation of the compound nucleus which has been investigated by measuring high-energy γ spectra for (isospin) symmetric and asymmetric reactions as in, e.g., [53; 54; 55]. Since in the extreme, such reactions essentially probe the di-nuclear system and not a compound nucleus (see, e.g., [56]), the resulting data are not entered into the present compilation. In general, in our compilations we focus more on low-energy data to avoid complications due to non-compound sources of high-energy γ rays, hence, data concerning the saturation or increase of the GDR width at very high excitation energies such as in [57; 58; 59] are typically omitted.

1.4. Data treatment

In our data treatment, the first step was to determine the target isotope from the context (for those few articles where it was not stated explicitly). Ranges of laboratory energy E_{lab} , initial excitation energy E_{ex} , and initial spin I_i were replaced by their central values. Average initial spins $\langle I_i \rangle$ were determined from maximum spins l_0 by means of $\langle I_i \rangle = \frac{2}{3}l_0$ for the case of fusion-evaporation reactions but irrespective of gating conditions. Ranges in final spin I_f after GDR γ emission were replaced by central values as well; the widths of the I_f ranges were converted into FWHM values which were preferred in this case over regular uncertainties due to the often non-Gaussian distribution of final spins.

Hot GDR parameterizations can take many different forms. The most common is probably the parameterization in terms of a centroid E , width Γ , and maximum σ of an equivalent Lorentzian photon-

absorption cross section. The maximum σ is often formulated as a fraction S of the Thomas-Reiche-Kuhn (TRK) sum rule which describes the integral $\frac{\pi}{2}\sigma\Gamma = 60\frac{NZ}{A}$ MeV mb of the Lorentzian in terms of neutron N , proton Z , and mass A numbers [15]. For two-component Lorentzian parameterizations, often the total $S_1 + S_2$, and either the ratio S_2/S_1 or the relative fraction $F_2 = S_2/(S_1 + S_2)$ are given. In either case, the given parameters were converted into S_1 and S_2 values. Sometimes, the total is assumed to fulfill the TRK while only the ratio or the relative fraction F_2 is determined by the fit. In these cases, the errors are marked by an asterisk to indicate correlations. In other cases, no information is given on the total. In those cases, the fraction S_2 is given in terms of the fraction S_1 in the table.

In case of GDR widths, some authors reduce their number of fit parameters by introducing a phenomenological relation between the widths and the centroids of a two-component Lorentzian according to $\Gamma = cE^2$, where c becomes the fit parameter [25]. In such cases, we have calculated the widths Γ including their errors. However, the errors are again correlated and marked by an asterisk. In the case of two GDR centroids, some authors give the average $E_{\text{ave}} = \frac{S_1 E_1 + S_2 E_2}{S_1 + S_2}$ and the ratio E_2/E_1 . Sometimes this ratio is replaced by an average deformation β which can be related to E_2/E_1 by either $\beta = \frac{2}{3}\sqrt{\frac{4\pi}{5}}\ln\frac{E_2}{E_1}$ (see, e.g., [60]) or $\beta = \sqrt{\frac{4\pi}{5}\frac{E_{\perp}/E_{\parallel}-1}{E_{\perp}/2E_{\parallel}+0.8665}}$ (see, e.g., [25]), where $E_{\perp}$ and $E_{\parallel}$ denote the centroids of the GDR components due to oscillations perpendicular and parallel to the symmetry axis. In the case where the original article did not mention which of the two formulas applies [61], the two formulas resulted in values for E_1 and E_2 within 0.2 MeV of each other, a difference far less than the quoted statistical error. In general, however, we did not concern ourselves with deformations. Only when oblate deformation was established from, e.g., the asymmetry a_2 , a '-' sign was added to the deformation parameter (in cases where the original article only provided the absolute value, see, e.g., [21]).

In our treatment of errors, the first step was to add quadratically statistical and systematic errors (when quoted separately). Where a range of systematic uncertainties is given, we adopted the center value of that range as a representative systematic uncertainty. When uncertainties are given in terms of a FWHM, it was converted by $\sigma = \text{FWHM}/\sqrt{8\ln 2}$. In general, rigorous error propagation was performed. However, no original work published the full covariance matrix for the fitted GDR parameters. Therefore, the derived errors are only representative of the true errors under the assumption that the originally fitted parameters are fully uncorrelated. In cases where we determined more parameters than were originally fitted, an asterisk denotes the correlations which were introduced to the errors. In cases where GDR parameters were held fixed during the fit, a little 'f' was added in the table instead of an error. Some works do not cite errors at all. In such cases we have made no attempt to estimate the errors. In a few cases where E_1 and E_2 were given without errors, while E_{ave} was given with error, we could not find a good method to translate this error into errors of the individual values, and hence E_1 and E_2 remain without errors (see, e.g., [21]).

1.5. Policy

Although we have tried to avoid a true evaluation of the original articles, we have excluded some of them from the present compilation. Typically, where high-energy γ rays from sources other than a compound nucleus were investigated (such as the mononucleus as in [29] or the di-nuclear system

as in [56]), the resulting data are not used for the present compilation. In a different case, the data in question were heavily contaminated by pile-up events which lead to unphysical GDR parameters [52]. In general, when different fits to the same data were performed (using different input parameters for the statistical-model calculation such as the level-density formula, see, e.g., [16], or the nuclear viscosity in the case of an open fission channel, see, e.g., [25]), we present all possible fits. On the other hand, when essentially the same data were presented in a conference proceedings as well as in a subsequent refereed article with no or minimal differences in the fit parameters, we typically report only the results from the refereed article with few exceptions. Finally, when no GDR parameters were given (see, e.g., [62; 63; 64] or when we feel that the statistical-model description of the experiment is rather tentative [65; 66], we made no attempt to fit the experimental spectrum ourselves, hence such data are not taken into account in the present compilation.

1.6. Note on References

When data from this compilation are cited, reference should also be made to the original publication as well.

Acknowledgments

We would like to thank Chris Kawatsu and Mark Shevin for the initial literature search. This work was supported by National Science Foundation Grant No. PHY01-10253.

REFERENCES

1. D. Brink, Ph.D. thesis, Oxford University, 1955.
2. P. Axel, Phys. Rev. **126**, 671 (1962).
3. P. d'Agostino, G. Fazio, G. Giardina, M. Herman, A. Lamberto, G.F. Rappazzo, R. Sturiale, F. Malaguti, D.O. Eremenko, N.V. Eremine, S.Yu. Platonov, and O.A. Yuminov, Bull. Russ. Acad. Scien. Phys. **62**, 1736 (1998).
4. Dimitri Kusnezov, Y. Alhassid, and K.A. Snover, Phys. Rev. Lett. **81**, 542 (1998); Dimitri Kusnezov, Y. Alhassid, K.A. Snover, and W.E. Ormand, Nucl. Phys. **A 687**, 212c (2001).
5. W.E. Ormand, P.F. Bortignon, and R.A. Broglia, Phys. Rev. Lett. **77**, 607 (1996).
6. Ahmad Ansari, Nguyen Dinh Dang, and Akita Arima, Phys. Rev. C **63**, 024310 (2001).
7. Yu.P. Popov, Fiz. Elem. Chastits At. Yadra **13**, 1165 (1982) [Sov. J. Part. Nucl. **13**, 483 (1982)].
8. S.G. Kadenskii, V.P. Markushev, and V.I. Furman, Yad. Fiz. **37**, 277 (1983) [Sov. J. Nucl. Phys. **37**, 165 (1983)].
9. V.K. Sirotkin, Yad. Fiz. **43**, 570 (1986) [Sov. J. Nucl. Phys. **43**, 362 (1986)].
10. J. Kopecky and M. Uhl, Phys. Rev. C **41**, 1941 (1990).

11. S.F. Mughabghab and C.L. Dunford, Phys. Lett. B **487**, 155 (2000).
12. O.T. Grudzevich, Yad. Fiz. **62**, 227 (1999) [Phys. At. Nucl. **62**, 192 (1999)].
13. A. Voinov, M. Guttormsen, E. Melby, J. Rekstad, A. Schiller, and S. Siem, Phys. Rev. C **63**, 044313 (2001).
14. V. Baran, M. Colonna, M. Di Toro, A. Guarnera, V.N Kondratyev, and A. Smerzi, Nucl. Phys. A **599**, 29c (1996).
15. Samuel S. Dietrich and Barry L. Berman, At. Data Nucl. Data Tables **38**, 199 (1998).
16. G. Feldman, K.A. Snover, J.A. Behr, C.A. Gossett, J.H. Gundlach, and M. Kicińska-Habior, Phys. Rev. C **47**, 1436 (1993).
17. S.K. Rathi, D.R. Chakrabarty, V.M. Datar, Suresh Kumar, E.T. Mirgule, A. Mitra, V. Nanal, and H.H. Oza, Phys. Rev. C **67**, 024603 (2003).
18. A. Bracco, J.J. Gaardhøje, A.M. Bruce, J.D. Garrett, B. Herskind, M. Pignanelli, D. Barnéoud, H. Nifenecker, J.A. Pinston, C. Ristori, F. Schussler, J. Bacelar, and H. Hofmann, Phys. Rev. Lett. **62**, 2080 (1989).
19. F. Lucarelli, N. Gelli, P. Blasi, M. Cinausero, E. Fioretto, G. Prete, D. Fabris, G. Nebbia, G. Viesti, E.M. Fiore, L. Fiore, V. Paticchio, and P.F. Bortignon, Z. Phys. A **355**, 35 (1996).
20. M.P. Kelly, J.F. Liang, A.A. Sonzogni, K.A. Snover, J.P.S. van Schagen, and J.P. Lestone, Phys. Rev. C **56**, 3201 (1997).
21. J.J. Gaardhøje, A.M. Bruce, and B. Herskind, Nucl. Phys. A **482**, 121c (1988).
22. M. Kmiecik, A. Maj, A. Bracco, F. Camera, B. Million, and O. Wieland, Eur. Phys. J. A **12**, 5 (2001).
23. A. Stolk, A. Balanda, M.N. Harakeh, W.H.A. Hesselink, J. Penninga, and H. Rijneveld, Nucl. Phys. A **505**, 241 (1989).
24. F. Camera, A. Bracco, V. Nanal, M.P. Carpenter, F. Della Vedova, S. Leoni, B. Million, S. Mantonani, M. Pignanelli, O. Wieland, B.B. Back, A.M. Heinz, R.V.F. Janssens, D. Jenkins, T.L. Khoo, F.G. Kondev, T. Lauritsen, C.F. Lister, B. McClintock, S. Mitsuoka, E.F. Moore, D. Seweryniak, R.H. Siemssen, R.J. Van Swol, D. Hofman, M. Thoennessen, K. Eisenman, P. Heckman, J. Seitz, R. Varner, M. Halbert, I. Dioszegi, and A. Lopez-Martens, Phys. Lett. B **560**, 155 (2003).
25. G. van 't Hof, J.C.S. Bacelar, I. Diószegi, M.N. Harakeh, W.H.A. Hesselink, N. Kalantar-Nayestanaki, A. Kugler, H. van der Ploeg, A.J.M. Plompen, and J.P.S. van Schagen, Nucl. Phys. A **638**, 613 (1998).
26. R. Butsch, M. Thoennessen, D.R. Chakrabarty, M.G. Herman, and P. Paul, Phys. Rev. C **41**, 1530 (1990).
27. F.S. Dietrich, J.C. Browne, W.J. O'Connell, and M.J. Kay, Phys. Rev. C **10**, 795 (1974).
28. N.P. Shaw, I. Diószegi, I. Mazumdar, A. Buda, C.R. Morton, J. Velkovska, J.R. Beene, D.W. Stracener, R.L. Varner, M. Thoennessen, and P. Paul, Phys. Rev. C **61**, 044612 (2000).
29. T.S. Tveter, J.J. Gaardhøje, A. Maj, T. Ramsøy, A. Ataç, J. Bacelar, A. Bracco, A. Buda, F. Camera, B. Herskind, W. Korten, W. Królas, A. Menthe, B. Million, H. Nifenecker, M. Pignanelli, J.A. Pinston, H.v.d. Ploeg, F. Schussler, and G. Sletten, Phys. Rev. Lett. **76**, 1035 (1996); T.S. Tveter, J.J. Gaardhøje, A. Maj, T. Ramsøy, A. Ataç, J. Bacelar, A. Bracco, A. Buda, F. Camera, B. Herskind,

- W. Korten, W. Królas, A. Menche, B. Million, H. Nifenecker, M. Pignanelli, J.A. Pinston, H. v. d. Ploeg, F. Schussler, and G. Sletten, Nucl. Phys. **A 599**, 123c (1996).
30. T. Baumann, E. Ramakrishnan, A. Azhari, J.R. Beene, R.J. Charity, J.F. Dempsey, M.L. Halbert, P.-F. Hua, R.A. Kryger, P.E. Mueller, R. Pfaff, D.G. Sarantites, L.G. Sobotka, D.W. Stracener, M. Thoennessen, G. Van Buren, R.L. Varner, and S. Yokoyama, Nucl. Phys. **A 635**, 428 (1998).
 31. G. Enders, F.D. Berg, K. Hagel, W. Kühn, V. Metag, R. Novotny, M. Pfeiffer, O. Schwalb, R.J. Charity, A. Gobbi, R. Freifelder, W. Henning, K.D. Hildebrand, R. Holzmann, R.S. Mayer, R.S. Simon, J.P. Wessels, G. Casini, A. Olmi, and A.A. Stefanini, Phys. Rev. Lett. **69**, 249 (1992).
 32. J.E. Draper, J.O. Newton, L.G. Sobotka, H. Lindenberger, G.J. Wozniak, L.G. Moretto, F.S. Stephens, R.M. Diamond, and R.J. McDonald, Phys. Rev. Lett. **49**, 434 (1982).
 33. N. Herrmann, R. Bock, H. Emling, R. Freifelder, A. Gobbi, E. Grosse, K.D. Hildebrand, R. Kulesa, T. Matulewicz, F. Rami, R.S. Simon, H. Stelzer, J. Wessels, P.R. Maurenzig, A. Olmi, A.A. Stefanini, W. Kühn, V. Metag, R. Novotny, M. Gnirs, D. Pelte, P. Braun-Munzinger, and L.G. Moretto, Phys. Rev. Lett. **60**, 1630 (1988).
 34. R. Hingmann, W. Kühn, V. Metag, R. Mühlhans, R. Novotny, A. Ruckelshausen, W. Cassing, H. Emling, R. Kulesa, H.J. Wollersheim, B. Haas, J.P. Vivien, A. Boullay, H. Delagrange, H. Doubre, C. Grégoire, and Y. Schutz, Phys. Rev. Lett. **58**, 759 (1987).
 35. M. Thoennessen, J.R. Beene, F.E. Bertrand, D.J. Horen, M.L. Halbert, D.C. Hensley, J.E. Lisantti, W. Mittig, Y. Schutz, N. Alamanos, F. Auger, J. Barrette, B. Fernandez, A. Gillibert, B. Haas, J.P. Vivien, and A.M. Nathan, Phys. Rev. C **43**, R12 (1991).
 36. A. Winther, Nucl. Phys. **A 594**, 203 (1995).
 37. W. Swiatecki, Nucl. Phys. **A 376**, 275 (1982).
 38. W. Hauser and H. Feshbach, Phys. Rev. **87**, 366 (1952).
 39. F. Pühlhofer, Nucl. Phys. **A 280**, 267 (1977).
 40. W. Reisdorf, Z. Phys. A **300**, 227 (1981).
 41. W. Dilg, W. Schantl, H. Vonach, and M. Uhl, Nucl. Phys. **A 217**, 269 (1973).
 42. A.V. Ignatyuk, K.K. Istekov, and G.N. Smirenkin, Yad. Fiz. **29**, 875 (1979) [Sov. J. Nucl. Phys. **29**, 450 (1979)].
 43. B.J. Fineman, K.-T. Brinkmann, A.L. Caraley, N. Gan, R.L. McGrath, and J. Velkovska, Phys. Rev. C **50**, 1991 (1994).
 44. W.D. Myers and W.J. Swiatecki, Nucl. Phys. **81**, 1 (1966).
 45. A. Bałanda, J.C.S. Bacelar, E. Běták, J.A. Bordewijk, A. Krasznahorka, H. van der Ploeg, R.H. Siemssen, H.W. Wilschut, and A. van der Woude, Nucl. Phys. **A 575**, 348 (1994).
 46. E. Běták, Report FU SAV 89/5, Bratislava 1989.
 47. M. Kicińska-Habior, K.A. Snover, J.A. Behr, C.A. Gossett, J.H. Gundelach, and G. Feldman, Phys. Rev. C **45**, 569 (1992).
 48. M. Kicińska-Habior, K.A. Snover, J.A. Behr, G. Feldman, C.A. Gossett, and J.H. Gundlach, Phys. Rev. C **41**, 2075 (1990).
 49. S. Flibotte, B. Haas, P. Taras, H.R. Andrews, D.C. Radford, and D. Ward, Nucl. Phys. **A 531**, 205

(1991).

50. J.J. Gaardhøje, C. Ellegaard, B. Herskind, R.M. Diamond, M.A. Deleplanque, G. Dines, A.O. Macchiavelli, and F.S. Stephens, *Phys. Rev. Lett.* **56**, 1783 (1986).
51. M. Kicińska-Habior, K.A. Snover, C.A. Gossett, J.A. Behr, G. Feldman, H.K. Glatzel, J.H. Gundlach, and E.F. Garman, *Phys. Rev. C* **36**, 612 (1987).
52. H.P. Morsch, W. Spang, J.R. Beene, F.E. Bertrand, R.L. Auble, J.L. Charvet, M.L. Halbert, D.C. Hensley, I.Y. Lee, R.L. Varner, D.G. Sarantites, and D.W. Stracener, *Phys. Rev. Lett.* **64**, 1999 (1990); M. Thoennessen, *Phys. Rev. Lett.* **66**, 1640 (1991); H.P. Morsch, W. Spang, J. Beene, and F. Bertrand, *Phys. Rev. Lett.* **66**, 1641 (1991).
53. D. Pierroutsakou, A. Boiano, A. De Rosa, M. Di Pietro, G. Inglima, M. La Commara, Ruhan Ming, B. Martin, R. Mordente, A. Ordine, F. Rizzo, V. Roca, M. Romoli, M. Sandoli, F. Soramel, L. Stroe, M. Trotta, and E. Vardaci, *Eur. Phys. J. A* **17**, 71 (2003).
54. M. Thoennessen, E. Ramakrishnan, J.R. Beene, F.E. Bertrand, M.L. Halbert, D.J. Horen, P.E. Mueller, and R.L. Varner, *Phys. Rev. C* **51**, 3148 (1995).
55. S. Flibotte, Ph. Chomaz, M. Colonna, M. Cromaz, J. DeGraaf, T.E. Drake, A. Galindo-Uribarri, V.P. Janzen, J. Jonkman, S.W. Marshall, S.M. Mullins, J.M. Nieminen, D.C. Radford, J.L. Rodriguez, J.C. Waddington, D. Ward, and J.N. Wilson, *Phys. Rev. Lett.* **77**, 1448 (1996).
56. L. Campajola, A. De Rosa, A. D’Onofrio, L. Gialanella, G. Inglima, M. La Commara, A. Ordine, D. Pierroutsakou, V. Roca, M. Romano, M. Romoli, M. Sandoli, F. Terrasi, G. Cardella, A. Di Pietro, A. Musumarra, M. Papa, G. Pappalardo, F. Rizzo, N. Alamanos, F. Auger, A. Gillibert, A. Incicchitti, D. Prosperi, G. De Angelis, E. Fioretto, and R. Bernabei, *Z. Phys. A* **352**, 421 (1995); L. Campajola, A. De Rosa, A. D’Onofrio, L. Gialanella, G. Inglima, M. La Commara, A. Ordine, D. Pierroutsakou, V. Roca, M. Romano, M. Romoli, M. Sandoli, F. Terrasi, F. Amorini, G. Cardella, R. Croce, A. Di Pietro, A. Musumarra, M. Papa, G. Pappalardo, F. Rizzo, J.P.S. Van Shagen, N. Alamanos, F. Auger, A. Gillibert, D. Prosperi, G. De Angelis, and E. Fioretto, *Nucl. Phys. A* **583**, 119c (1995); M. Sandoli, L. Campajola, A. De Rosa, A. D’Onofrio, G. Inglima, M. La Commara, A. Ordine, D. Pierroutsakou, V. Roca, M. Romano, M. Romoli, F. Terrasi, M. Trotta, G. Cardella, M. Papa, G. Pappalardo, F. Rizzo, N. Alamanos, F. Auger, and A. Gillibert, *Z. Phys. A* **357**, 67 (1997); D. Pierroutsakou, A. Boiano, A. De Rosa, M. Di Pietro, G. Inglima, M. La Commara, Ruhan Ming, B. Martin, R. Mordente, A. Ordine, F. Rizzo, V. Roca, M. Romoli, M. Sandoli, F. Soramel, L. Stroe, M. Trotta, and E. Vardaci, *Eur. Phys. J. A* **17**, 71 (2003).
57. K. Yoshida, J. Kasagi, H. Hama, M. Sakurai, M. Kodama, K. Furutaka, K. Ieki, W. Galster, T. Kubo, and M. Ishihara, *Phys. Lett. B* **245**, 7 (1990); J. Kasagi, K. Furutaka, T. Murakami, A. Yajima, M. Ohshima, S. Niiya, H. Tominaga, K. Yoshida, H. Hama, K. Ieki, W. Galster, K. Kubo, M. Ishihara, and A. Galonsky, *Nucl. Phys. A* **538**, 585c (1992).
58. J.H. Le Faou, T. Suomijärvi, Y. Blumentfeld, P. Piattelli, C. Agodi, N. Alamanos, R. Alba, F. Auger, G. Bellia, Ph. Chomaz, R. Coniglione, A. Del Zoppo, P. Finocchiaro, N. Frascaria, J.J. Gaardhøje, J.P. Garron, A. Gillibert, M. Laméhi-Rachti, R. Liguori-Neto, C. Maiolino, E. Migneco, G. Russo, J.C. Roynette, D. Santonocito, P. Sapienza, J.A. Scarpaci, and A. Smerzi, *Phys. Rev. Lett.* **72**, 3321 (1994); T. Suomijärvi, Y. Blumenfeld, P. Piattelli, J.H. Le Faou, C. Agodi, N. Alamanos, R. Alba, F. Auger, G. Bellia, Ph. Chomaz, R. Coniglione, A. Del Zoppo, P. Finocchiaro, N. Frascaria, J.J. Gaardhøje, J.P. Garron, A. Gillibert, M. Laméhi-Rachti, R. Liguori-Neto, C. Maiolino, E. Migneco,

- G. Russo, J.C. Roynette, D. Santonocito, P. Sapienza, J.A. Scarpaci, and A. Smerzi, *Phys. Rev. C* **53**, 2258 (1996); T. Suomijärvi, J.H. Le Faou, Y. Blumenfeld, P. Piattelli, C. Agodi, N. Alamanos, R. Alba, F. Auger, G. Bellia, Ph. Chomaz, R. Coniglione, A. Del Zoppo, P. Finocchiaro, N. Frascaria, J.J. Gaardhøje, J.P. Garron, A. Gillibert, M. Laméhi-Rachti, R. Liguori-Neto, K. Loukachine, C. Maiolino, E. Migneco, S. Montironi, G. Russo, J.C. Roynette, D. Santonocito, P. Sapienza, J.A. Scarpaci, and A. Smerzi, *Nucl. Phys. A* **583**, 105 (1995).
59. J.J. Gaardhøje, A.M. Bruce, J.D. Garrett, B. Herskind, D. Barnéoud, M. Maurel, H. Nifenecker, J.A. Pinston, P. Perrin, C. Ristori, F. Schussler, A. Bracco, and M. Pignanelli, *Phys. Rev. Lett.* **59**, 1409 (1987).
 60. M. Kicińska-Habior, K.A. Snover, J.A. Behr, C.A. Gossett, Y. Alhassid, and N. Whelan, *Phys. Lett. B* **308**, 225 (1990).
 61. P. Thirolf, D. Habs, D. Schwalm, R.D. Fischer, and V. Metag, *Nucl. Phys. A* **482**, 93c (1988).
 62. W. Hennerici, V. Metag, H.J. Hennrich, R. Repnow, W. Wahl, D. Habs, K. Helmer, U. v. Helmolt, H.W. Heyng, B. Kolb, D. Pelte, D. Schwalm, R.S. Simon, and R. Albrecht, *Nucl. Phys. A* **396**, 329c (1983).
 63. L.H. Zhu, D. Bazzacco, S. Lunardi, G. Viesti, M. Cinausero, G. de Angelis, M. De Poli, A. Gadea, D.R. Napoli, C.M. Petrache, and C. Rossi Alvarez, *Phys. Rev. C* **55**, 1169 (1997).
 64. Adam Maj, Maria Kmiecik, Nicolas Schunk, and Jan Styczén, *AIP Conf. Proc.* **802**, 264 (2005).
 65. J.O. Newton, B. Herskind, R.M. Diamond, E.L. Dines, J.E. Draper, K.H. Lindenberg, C. Schück, S. Shih, and F.S. Stephens, *Phys. Rev. Lett.* **46**, 1383 (1981).
 66. J.P.S. van Schagen, F. Amorini, G. Cardella, R. Croce, A. DiPietro, A. Musumarra, M. Papa, G. Pappalardo, F. Rizzo, L. Campajola, A. DeRosa, A. D’Onofrio, G. Inghima, M. LaCommara, D. Pierroutsakou, V. Rocca, M. Pomano, M. Romoli, M. Sandoli, and F. Terrasi, *Nucl. Phys. A* **583**, 123c (1995).

EXPLANATION OF TABLE

TABLE. Reaction and GDR parameters

No.	number of item
Ref.	reference
Pro.	projectile nucleus
Tar.	target nucleus
CN	compound nucleus as determined by the gating conditions
E_{lab}	laboratory beam energy (MeV), sometimes given at the center of the target
E_{ex}	excitation energy of the compound nucleus (MeV) as determined by the gating conditions, sometimes averaged over the target thickness
$\langle I_i \rangle$	initial average angular momentum of the compound nucleus ($\hbar$) as determined by the gating conditions, sometimes calculated by $\langle I_i \rangle = \frac{2}{3}l_0$ where l_0 is the maximum spin of the compound system calculated by the theories of Winther or Swiatecki
T_i	initial temperature of the compound nucleus (MeV) as determined by the gating conditions; sometimes determined before or after pre-equilibrium particle emission, but always before particle evaporation
l.d.	level density parameterization: R=Reisdorf, P=Pühlhofer, I=Ignatyuk, D=Dilg, F=Fine-man; in some cases, the Wigner energy is included in the parameterization
A/a	Level density parameter (MeV), often used only for conversion of excitation energy to temperature of the compound nucleus; the default value for the Pühlhofer l.d. parameterization is $A/a = 8$ MeV which should be assumed if no value is given
$\langle I_f \rangle$	final average angular momentum of the compound nucleus ($\hbar$) after high-energy γ emission and influenced by gating conditions
FWHM	full width at half maximum of the final angular momentum distribution ($\hbar$)
E_f	final average excitation energy of the compound nucleus (MeV) after high-energy γ emission and influenced by gating conditions
T_f	final temperature of the compound nucleus (MeV) after high-energy γ emission and influenced by gating conditions
Code	statistical model computer code used for fitting the experimental spectra: C=Cascade (many different modifications exist), P=Pace, P2=Peqag2; if no code is given, most likely some version of the code Cascade was used
gate	gating condition: $\alpha d = \alpha$ decay, $\gamma c = \gamma$ coincidences, $d\gamma t =$ discrete γ transitions, $did =$ discrete isomeric decay, $dis =$ deep inelastic scattering, $er =$ evaporation residues, $f = \gamma$ -fold, $ff =$ fission fragments, $ipc =$ internal pair conversion, $is =$ inelastic scattering, $lcp =$ light charged particle, $\gamma m = \gamma$ multiplicity, $\gamma \Sigma E = \gamma$ sum energy, $st =$ subtraction technique
S_1	strength of the first component of the GDR (fraction of the TRK sum rule)
E_1	centroid of the first component of the GDR (MeV)
Γ_1	width of the first component of the GDR (MeV)
S_1	strength of the second component of the GDR (fraction of the TRK sum rule)
E_1	centroid of the second component of the GDR (MeV)
Γ_1	width of the second component of the GDR (MeV)
FWHM	full width at half maximum of the GDR cross-section parameterization (MeV)

def. average deformation β or in some cases δ of the compound nucleus after high-energy γ emission

Throughout the table, *italics* entries refer to derived values, i.e., anything which is not found verbatim in the cited reference. A little 'f' instead of an uncertainty means that the value in question was kept fixed in the fit. An asterisk behind any two given uncertainties means that these uncertainties are correlated by the specific parameterization used during fitting. For entries 48 and 52, nuclear bremsstrahlung was given as a source of systematic error, however, it was not quantified in the original article. Entries 245, 248, 252, and 255 fitted the GDR using a triaxial parameterization. In order to reduce the number of free parameters, they chose $S_1 = S_2 = S_3 = 1/3$ and $\Gamma_1 = \Gamma_2 = \Gamma_3 = \Gamma$. The given fit parameters are for those entries are: Γ and the three centroids E_1 , E_2 , and E_3 from which the two deformation parameters β and γ were deduced. Entry 274 employed a Hill-Wheeler parameterization [4] and takes the temperature dependence of the GDR into account already in the statistical-model calculation instead of giving an averaged parameterization of the GDR over the investigated energy and spin range. Hence, it gives only a spherical width and how this width scales with the GDR centroid energy. Entries 288 and 289 belong to one fit, where different parameters were employed above and below $\langle I_f \rangle = 14\hbar$.

TABLE I. Giant Electric Dipole Resonance parameters.

See page 13 for Explanation of Tables.

This space intentionally left blank

No.	Ref.	Pro.	Tar.	CN	E_{lab}	E_{ex}	$\langle I_i \rangle$	T_i	l.d.	A/a	$\langle I_f \rangle$	FWHM	E_f	T_f	Code	gate
1	Kic90	¹² C	²⁷ Al	³⁹ K	33.8	40.0	7.7		R	8	8.0		14.9	1.75	C	
2	Kic90	¹² C	²⁷ Al	³⁹ K	48.2	50.0	13.2		R	8	13.0		14.3	1.71	C	
3	Kic90	¹² C	²⁷ Al	³⁹ K	62.7	60.0	16.5		R	8	16.0		13.9	1.69	C	
4	Kic90	¹³ C	²⁷ Al	⁴⁰ K	39.7	46.3	9.3		R	8	9.5		17.8	1.88	C	
5	Kic90	¹⁵ N	²⁷ Al	⁴² Ca	29.2	40.2	7.7		R	8	8.0		16.5	1.77	C	
6	Kic90	¹⁵ N	²⁷ Al	⁴² Ca	44.7	50.2	12.5		R	8	12.5		16.7	1.78	C	
7	Kic90	¹⁸ O	²⁷ Al	⁴⁵ Sc	31.5	42.0	7.7		R	8	8.0		16.1	1.69	C	
8	Kic90	¹⁸ O	²⁷ Al	⁴⁵ Sc	44.9	50.0	13.2		R	8	13.0		16.8	1.73	C	
9	Kic93	¹⁸ O	²⁷ Al	⁴⁵ Sc	44.9	50.0	13		R		13			1.7	C	
10	Kic90	¹⁸ O	²⁷ Al	⁴⁵ Sc	61.5	60.0	16.3		R	8	16.0		17.7	1.77	C	
11	Kic90	¹⁸ O	²⁷ Al	⁴⁵ Sc	72.5	66.6	18.7		R	8	18.5		17.8	1.78	C	
12	Kic93	¹⁸ O	²⁷ Al	⁴⁵ Sc	72.5	66.6	19.0		R		18.5			1.8	C	
13	Kic93	¹⁸ O	²⁷ Al	⁴⁵ Sc	89.9	76.7	22		R		21.4			2.0	C	
14	Kic93	¹⁸ O	²⁷ Al	⁴⁵ Sc	109.6	88.9	24		R		23.5			2.3	C	
15	Fel93	¹⁹ F	²⁷ Al	⁴⁶ Ti	29.0	42.5	3.5		R		3.4			1.93	C	
16	Fel93	¹⁹ F	²⁷ Al	⁴⁶ Ti	29.0	42.5	3.5		P	8.5	3.4			1.93	C	
17	Fel93	¹⁹ F	²⁷ Al	⁴⁶ Ti	36.4	46.8	8.6		R		8.2			1.85	C	
18	Fel93	¹⁹ F	²⁷ Al	⁴⁶ Ti	36.4	46.8	8.6		P	8.5	8.2			1.85	C	
19	Fel93	¹⁹ F	²⁷ Al	⁴⁶ Ti	45.3	52.0	14.3		R		13.9			1.67	C	
20	Fel93	¹⁹ F	²⁷ Al	⁴⁶ Ti	45.3	52.0	14.3		P	8.5	13.9			1.67	C	
21	Fel93	¹⁹ F	²⁷ Al	⁴⁶ Ti	55.7	58.1	19.1		R		18.2			1.50	C	
22	Fel93	¹⁹ F	²⁷ Al	⁴⁶ Ti	55.7	58.1	19.1		P	8.5	18.2			1.50	C	
23	Fel93	⁴ He	⁴⁸ Ti	⁵² Cr	12.0	20.4	4.5		R		4.5			0.51	C	
24	Fel93	⁴ He	⁴⁸ Ti	⁵² Cr	12.0	20.4	4.5		P	8.5	4.5			0.51	C	
25	Fel93	⁴ He	⁴⁸ Ti	⁵² Cr	17.2	25.2	6.5		R		6.5			0.89	C	
26	Fel93	⁴ He	⁴⁸ Ti	⁵² Cr	17.2	25.2	6.5		P	8.5	6.5			0.89	C	
27	Fel93	⁴ He	⁴⁸ Ti	⁵² Cr	24.0	31.5	8.4		R		8.4			1.26	C	
28	Fel93	⁴ He	⁴⁸ Ti	⁵² Cr	24.0	31.5	8.4		P	8.5	8.4			1.26	C	
29	Fel93	⁴ He	⁴⁸ Ti	⁵² Cr	28.0	35.2	9.3		R		9.3			1.36	C	
30	Fel93	⁴ He	⁴⁸ Ti	⁵² Cr	28.0	35.2	9.3		P	8.5	9.3			1.36	C	
31	Kro92	⁴ He	⁵¹ V	⁵⁵ Mn	26	31	8		P				8	1.1	C	
32	Vie89	³² S	²⁷ Al	⁵⁹ Cu	89	54.0	17		P						C	
33	Vie89	³² S	²⁷ Al	⁵⁹ Cu	89	54.0	17		P						C	
34	Vie89	³² S	²⁷ Al	⁵⁹ Cu	89	54.0	17		P						C	
35	Dre95	³² S	²⁷ Al	⁵⁹ Cu	90	54.4	15.0		R	8	12.8			1.7	C	
36	Dre95	³² S	²⁷ Al	⁵⁹ Cu	120	68.1	22.4		R	8	17.6			1.8	C	
37	Vie89	³² S	²⁷ Al	⁵⁹ Cu	140	77.4	25		P						C	
38	Vie89	³² S	²⁷ Al	⁵⁹ Cu	140	77.4	25		P						C	
39	Vie89	³² S	²⁷ Al	⁵⁹ Cu	140	77.4	25		P						C	
40	Dre95	³² S	²⁷ Al	⁵⁹ Cu	141	77.7	26.5		R	8	22.0			1.9	C	
41	Dre95	³² S	²⁷ Al	⁵⁹ Cu	175	93.2	28.7		R	8	27.5			2.0	C	$f \geq 2$
42	Dre95	³² S	²⁷ Al	⁵⁹ Cu	175	93.2	28.7		R	8	27.5			2.0	C	
43	Dre95	³² S	²⁷ Al	⁵⁹ Cu	215	111.4	30.7		R	8	31.5			2.1	C	$f \geq 2$
44	Dre95	³² S	²⁷ Al	⁵⁹ Cu	215	111.4	30.7		R	8	31.5			2.1	C	
45	Kic87	⁴ He	⁵⁹ Co	⁶³ Cu	17.8	22.5	4.5		R	8	5.5		4.3	0.74	C	
46	Kic87	⁴ He	⁵⁹ Co	⁶³ Cu	23.9	28.1	7.5		R	8	7.8		7.3	0.96	C	
47	Kic87	⁶ Li	⁵⁷ Fe	⁶³ Cu	13.9	32.0	2.0		R	8	3.0		12.0	1.23	C	
48	Kic87	⁴ He	⁵⁹ Co	⁶³ Cu	28.4	32.4	5.5		R	8	7.5		11.1	1.19	C	
49	Kic87	¹² C	⁵¹ V	⁶³ Cu	26.3	34.7	3.0		R	8	4.5		12.9	1.28	C	
50	Kic87	⁶ Li	⁵⁷ Fe	⁶³ Cu	22.1	39.5	7.0		R	8	7.5		17.7	1.50	C	
51	Kic87	¹² C	⁵¹ V	⁶³ Cu	32.2	39.5	8.5		R	8	9.5		16.6	1.45	C	
52	Kic87	⁶ Li	⁵⁷ Fe	⁶³ Cu	36.0	52.0	11.5		R	8	10.0		23.7	1.73	C	
53	Kic87	¹² C	⁵¹ V	⁶³ Cu	48.0	52.2	15.5		R	8	16.0		19.2	1.56	C	
54	Kic87	¹⁸ O	⁴⁵ Sc	⁶³ Cu	39.9	52.2	8.5		R	8	8.5		24.6	1.77	C	
55	Gar83	¹² C	⁵¹ V	⁶³ Cu		54.3	16								C	
56	Gar83	¹⁸ O	⁴⁵ Sc	⁶³ Cu		54.3	10								C	
57	Dre95	¹⁸ O	⁴⁵ Sc	⁶³ Cu	52	61.1	17.3		R	8	14.9			1.8	C	

No.	S_1	E_1	Γ_1	S_2	E_2	Γ_2	FWHM	def.
1	0.89 ± 0.17	16.4 ± 0.5	10.5 ± 1.0					
2	0.77 ± 0.13	16.8 ± 0.5	11.6 ± 1.1					
3	0.86 ± 0.14	17.0 ± 0.5	13.4 ± 1.1					
4	1.33 ± 0.22	17.2 ± 0.5	12.1 ± 1.1					
5	1.19 ± 0.20	17.7 ± 0.5	11.6 ± 1.0					
6	1.17 ± 0.19	17.1 ± 0.5	11.9 ± 1.0					
7	0.75 ± 0.12	17.0 ± 0.5	11.6 ± 1.0					
8	0.76 ± 0.12	16.1 ± 0.5	12.1 ± 1.1					
9	$0.86^{+0.14}_{-0.49}$	$16.3^{+1.8}_{-3.5}$	10.7 ± 1.9	$0.10^{+0.48}_{-0.09}$	$22.6^{+3.7}_{-5.6}$	9.9 ± 2.6	12.2 ± 0.4	0.35 ± 0.17
10	0.87 ± 0.14	16.1 ± 0.5	12.4 ± 1.1					
11	0.97 ± 0.16	16.3 ± 0.5	14.7 ± 1.1					
12	$0.83^{+0.05}_{-0.10}$	$16.0^{+0.5}_{-0.8}$	10.2 ± 0.8	$0.11^{+0.10}_{-0.04}$	$23.8^{+0.9}_{-1.4}$	5.5 ± 3.5	13.5 ± 0.5	0.42 ± 0.03
13	$0.94^{+0.12}_{-0.46}$	$16.3^{+1.0}_{-3.1}$	11.2 ± 1.2	$0.24^{+0.45}_{-0.11}$	$25.0^{+1.7}_{-4.8}$	8.3 ± 4.4	16.1 ± 0.7	0.45 ± 0.04
14	$0.97^{+0.18}_{-1.37}$	$15.9^{+1.5}_{-8.2}$	12.7 ± 2.2	$0.32^{+1.36}_{-0.17}$	$24.7^{+2.5}_{-12.8}$	11.0 ± 6.8	18.1 ± 0.9	0.46 ± 0.05
15	1.21 ± 0.14	18.0 ± 0.6	12.6 ± 2.0					
16	1.09 ± 0.15	16.9 ± 0.8	12.0 ± 2.0					
17	1.01 ± 0.03	17.7 ± 0.1	10.0 ± 0.3					
18	0.86 ± 0.04	17.1 ± 0.3	9.5 ± 0.3					
19	1.10 ± 0.03	17.9 ± 0.1	11.4 ± 0.3					
20	0.88 ± 0.06	17.1 ± 0.4	10.9 ± 0.3					
21	1.17 ± 0.02	18.0 ± 0.1	12.8 ± 0.2					
22	0.88 ± 0.06	16.8 ± 0.4	12.2 ± 0.5					
23	1.10 ± 0.15	17.7 ± 0.5	6.0 ± 1.0					
24	0.75 ± 0.15	17.7 ± 0.5	7.0 ± 1.0					
25	1.02 ± 0.05	17.6 ± 0.2	5.9 ± 0.3					
26	0.70 ± 0.04	17.9 ± 0.2	6.9 ± 0.4					
27	0.94 ± 0.03	18.1 ± 0.1	5.6 ± 0.2					
28	0.84 ± 0.10	17.7 ± 0.5	8.1 ± 0.5					
29	1.04 ± 0.03	18.7 ± 0.1	6.5 ± 0.2					
30	1.04 ± 0.16	17.9 ± 0.7	8.2 ± 0.5					
31	0.95 ± 0.02	17.4 ± 0.1	9.5 ± 0.5					
32	1.37 ± 0.04	17.5 ± 0.2	9.3 ± 0.4					
33	1.21 ± 0.04	17.9 ± 0.4	9.2 ± 0.7					
34	0.96 ± 0.12	17.1 ± 0.4	9.6 ± 1.3					
35	0.9 ± 0.1	17.9 ± 0.4	9.1 ± 0.6					
36	0.9 ± 0.1	17.6 ± 0.3	9.7 ± 0.6					
37	0.93 ± 0.20	16.6 ± 0.8	15.0 ± 3.1					
38	0.64 ± 0.11	17.8 ± 1.1	16.7 ± 1.1					
39	0.82 ± 0.11	15.7 ± 2.6	15.3 ± 0.7					
40	1.0 ± 0.1	17.9 ± 0.3	11.7 ± 0.6					
41	1.1 ± 0.1	17.6 ± 0.4	13.5 ± 0.6					
42	1.2 ± 0.2	17.4 ± 0.3	13.6 ± 0.7					
43	1.1 ± 0.1	17.3 ± 0.4	13.4 ± 1.0					
44	1.1 ± 0.2	17.1 ± 0.3	14.4 ± 0.8					
45	1.25 ± 0.08	17.1 ± 0.1	7.8 ± 0.6					
46	0.87 ± 0.08	17.4 ± 0.4	7.5 ± 0.4					
47	0.74 ± 0.10	16.5 ± 0.6	8.2 ± 0.8					
48	2.09 ± 0.20	17.2 ± 1.2	7.9 ± 0.3	brems				
49	1.37 ± 0.12	17.1 ± 0.5	8.1 ± 0.6					
50	0.80 ± 0.07	17.1 ± 0.5	8.8 ± 0.3					
51	0.97 ± 0.05	16.7 ± 0.3	7.7 ± 0.2					
52	0.87 ± 0.07	17.6 ± 0.4	10.2 ± 0.3	brems				
53	0.87 ± 0.07	16.8 ± 0.3	9.3 ± 0.3					
54	1.10 ± 0.07	16.5 ± 0.4	9.6 ± 0.4					
55	1.3 f	15.7 ± 0.63	7.5 ± 0.63					
56	1.3 f	15.7 ± 0.63	7.5 ± 0.63					
57	1.1 ± 0.2	17.3 ± 0.3	11.1 ± 0.6					

No.	Ref.	Pro.	Tar.	CN	E_{lab}	E_{ex}	$\langle I_i \rangle$	T_i	l.d.	A/a	$\langle I_f \rangle$	FWHM	E_f	T_f	Code	gate
58	Kic87	^{18}O	^{45}Sc	^{63}Cu	53.1	61.6	15.5		R	8	15.0		27.1	1.86	C	
59	Kic87	^{12}C	^{51}V	^{63}Cu	64.7	65.7	20.0		R	8	20.5		23.2	1.72	C	
60	Dre95	^{18}O	^{45}Sc	^{63}Cu	68	72.4	23.3		R	8	19.6			1.9	C	
61	Kic87	^{18}O	^{45}Sc	^{63}Cu	75.1	77.4	23.0		R	8	22.0		29.2	1.93	C	
62	Dre95	^{18}O	^{45}Sc	^{63}Cu	105	99.1	25.3		R	8	22.6			2.2	C	
63	Dre95	^{18}O	^{45}Sc	^{63}Cu	125	113.0	25.3		R	8	22.9			2.3	C	
64	Dre95	^{18}O	^{45}Sc	^{63}Cu	149	130.1	25.3		R	8	22.9			2.5	C	
65	Gar83	^{12}C	^{64}Zn	^{76}Kr		44.0	15								C	
66	Gar83	^{18}O	^{58}Ni	^{76}Kr		44.0	9								C	
67	Gar83	^{12}C	^{64}Zn	^{76}Kr		46.3	15								C	
68	Gar83	^{18}O	^{58}Ni	^{76}Kr		46.3	12								C	
69	Gar83	^{12}C	^{64}Zn	^{76}Kr		53.9	19								C	
70	Gar83	^{18}O	^{58}Ni	^{76}Kr		53.9	17								C	
71	Gun90	^{18}O	^{72}Ge	^{90}Zr	50	54.4					9			1.6	C	
72	Gun90	^{18}O	^{72}Ge	^{90}Zr	74	75.0					22			1.7	C	
73	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	12.4	14.1		1.28 ± 0.27	C	f=2-3
74	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	10.8	13.0		1.30 ± 0.27	C	f=2-3
75	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	15.0	15.1		1.25 ± 0.26	C	f=3-4
76	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	13.5	14.6		1.27 ± 0.26	C	f=3-4
77	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	17.5	14.6		1.21 ± 0.26	C	f=4-5
78	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	16.0	14.6		1.23 ± 0.26	C	f=4-5
79	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	20.0	14.1		1.17 ± 0.26	C	f=5-6
80	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	18.7	14.1		1.19 ± 0.26	C	f=5-6
81	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	22.0	13.0		1.13 ± 0.26	C	f=6-7
82	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	20.9	13.0		1.15 ± 0.26	C	f=6-7
83	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	23.6	12.0		1.09 ± 0.23	C	f=7-14
84	Rat03	^{28}Si	^{58}Ni	^{86}Mo	100	49	20		I	7.5	23.1	12.0		1.10 ± 0.23	C	f=7-14
85	Rat03	^{28}Si	^{58}Ni	^{86}Mo	125	66	31		I	7.5	10.6	14.1		1.60 ± 0.34	C	f=1-2
86	Rat03	^{28}Si	^{58}Ni	^{86}Mo	125	66	31		I	7.5	13.3	17.0		1.57 ± 0.34	C	f=2-3
87	Rat03	^{28}Si	^{58}Ni	^{86}Mo	125	66	31		I	7.5	16.8	19.1		1.53 ± 0.34	C	f=3-4
88	Rat03	^{28}Si	^{58}Ni	^{86}Mo	125	66	31		I	7.5	20.6	20.0		1.47 ± 0.34	C	f=4-5
89	Rat03	^{28}Si	^{58}Ni	^{86}Mo	125	66	31		I	7.5	24.1	19.3		1.41 ± 0.33	C	f=5-6
90	Rat03	^{28}Si	^{58}Ni	^{86}Mo	125	66	31		I	7.5	26.9	18.1		1.36 ± 0.31	C	f=6-7
91	Rat03	^{28}Si	^{58}Ni	^{86}Mo	125	66	31		I	7.5	29.2	17.2		1.31 ± 0.31	C	f=7-8
92	Rat03	^{28}Si	^{58}Ni	^{86}Mo	125	66	31		I	7.5	31.2	16.0		1.26 ± 0.30	C	f=8-14
93	Kic92	^{16}O	^{76}Se	^{92}Mo	50.0	48.1	9.0		R	8	9.0			1.35	C	
94	Kic92	^{16}O	^{76}Se	^{92}Mo	72.2	66.5	19.7		R	8	19.5			1.45	C	
95	Gun90	^{28}Si	^{64}Ni	^{92}Mo	137	93.5					33			2.0	C	
96	Kic92	^{18}O	^{82}Se	^{100}Mo	49.1	48.1	9.3		R	8	9.3			1.35	C	
97	Kic92	^{18}O	^{82}Se	^{100}Mo	63.4	59.8	19.7		R	8	19.5			1.45	C	
98	Kic92	^{18}O	^{82}Se	^{100}Mo	72.8	67.5	24.2		R	8	24.0			1.50	C	
99	Luc96	^{32}S	^{74}Ge	^{106}Cd	210	134	46			7.5					C	lcp
100	Kel97	^{18}O	^{100}Mo	^{115}In	200	137			R						C	lcp
101	Mat97	^{58}Ni	^{48}Ti	^{106}Sn	260	80					24				C	f=3-6 & er
102	Mat97	^{58}Ni	^{48}Ti	^{106}Sn	260	80					36				C	f=7-20 & er
103	Gaa84	^{12}C	^{96}Ru	^{108}Sn	63	51.0				7.5					C	γ_c
104	Gaa84	^{16}O	^{92}Mo	^{108}Sn	73	51.8				7.5					C	γ_c
105	Gaa88	^{16}O	^{92}Mo	^{108}Sn		57				8	12					$\gamma_{\Sigma E}$
106	Gaa88	^{16}O	^{92}Mo	^{108}Sn		57				8	12					$\gamma_{\Sigma E}$
107	Gaa88	^{16}O	^{92}Mo	^{108}Sn		57				8	20					$\gamma_{\Sigma E}$
108	Gaa88	^{16}O	^{92}Mo	^{108}Sn		57				8	26					$\gamma_{\Sigma E}$
109	Gaa84	^{16}O	^{92}Mo	^{108}Sn	84	61.2				7.5					C	γ_c
110	Bra95	^{48}Ti	^{61}Ni	^{109}Sn	203	80				8	45	24			C	f>9
111	Bra95	^{48}Ti	^{61}Ni	^{109}Sn	203	80		1.79		8	40	18		1.6 ± 0.13	C	γ_m
112	Bra95	^{48}Ti	^{61}Ni	^{109}Sn	203	80		1.57		8	49	16		1.4 ± 0.13	C	γ_m
113	Tho95	^{18}O	^{92}Mo	^{110}Sn	72	56				10	22.5	15			C	f=5-10
114	Tho95	^{50}Ti	^{60}Ni	^{110}Sn	163	56				10	22.5	15			C	f=5-10

No.	S_1	E_1	Γ_1	S_2	E_2	Γ_2	FWHM	def.
58	0.99 ± 0.10	16.3 ± 0.3	9.9 ± 0.5					
59	1.07 ± 0.10	17.0 ± 0.3	10.5 ± 0.5					
60	0.9 ± 0.1	17.3 ± 0.3	11.4 ± 0.6					
61	0.90 ± 0.10	16.4 ± 0.3	10.6 ± 0.6					
62	1.1 ± 0.1	16.8 ± 0.3	14.8 ± 0.9					
63	1.1 ± 0.2	16.7 ± 0.3	12.7 ± 2.3					
64	1.2 ± 0.2	16.5 ± 0.3	13.0 ± 2.1					
65	1.3 f	15.5 ± 0.63	7.0 ± 0.63					
66	1.3 f	15.5 ± 0.63	7.0 ± 0.63					
67	1.3 f	15.5 ± 0.63	7.0 ± 0.63					
68	1.3 f	15.5 ± 0.63	7.0 ± 0.63					
69	1.3 f	15.5 ± 0.63	7.0 ± 0.63					
70	1.3 f	15.5 ± 0.63	7.0 ± 0.63					
71	0.26 ± 0.15	14.6 ± 0.7	3.42 ± 0.80	0.86 ± 0.18	17.7 ± 0.5	7.02 ± 1.25	8.22 ± 0.24	
72	0.36 ± 0.13	14.6 ± 0.5	5.24 ± 0.88	0.68 ± 0.14	18.2 ± 0.6	9.07 ± 0.77	9.67 ± 0.39	
73	0.33 f	14.6 ± 0.3	4.5 ± 0.3	0.67 f	18.9 ± 0.6	6.3 ± 0.7	8.8 ± 0.5	0.31 ± 0.05
74	0.33 f	14.4 ± 0.4	4.0 ± 0.4	0.67 f	18.7 ± 0.9	6.0 ± 0.9	8.5 ± 0.8	0.31 ± 0.06
75	0.33 f	15.0 ± 0.3	4.5 ± 0.3	0.67 f	18.9 ± 0.5	5.7 ± 0.5	8.0 ± 0.4	0.28 ± 0.04
76	0.33 f	14.8 ± 0.4	4.2 ± 0.4	0.67 f	18.9 ± 0.8	5.1 ± 0.7	7.7 ± 0.7	0.29 ± 0.05
77	0.33 f	15.2 ± 0.3	4.0 ± 0.3	0.67 f	19.0 ± 0.5	5.2 ± 0.5	7.5 ± 0.4	0.27 ± 0.04
78	0.33 f	14.9 ± 0.4	4.0 ± 0.4	0.67 f	18.7 ± 0.5	4.8 ± 0.6	7.2 ± 0.5	0.27 ± 0.05
79	0.33 f	14.9 ± 0.3	3.5 ± 0.3	0.67 f	18.5 ± 0.4	4.4 ± 0.4	6.7 ± 0.4	0.26 ± 0.04
80	0.33 f	15.0 ± 0.4	3.5 ± 0.4	0.67 f	18.4 ± 0.5	4.5 ± 0.6	6.6 ± 0.4	0.24 ± 0.05
81	0.33 f	14.5 ± 0.3	3.3 ± 0.3	0.67 f	17.7 ± 0.4	3.6 ± 0.4	5.7 ± 0.3	0.24 ± 0.04
82	0.33 f	14.5 ± 0.4	3.2 ± 0.4	0.67 f	17.8 ± 0.5	3.4 ± 0.6	5.7 ± 0.4	0.24 ± 0.05
83	0.33 f	14.2 ± 0.3	3.0 ± 0.3	0.67 f	17.7 ± 0.5	3.9 ± 0.5	6.2 ± 0.4	0.26 ± 0.04
84	0.33 f	14.5 ± 0.4	2.9 ± 0.4	0.67 f	17.4 ± 0.5	3.4 ± 0.6	5.3 ± 0.4	0.22 ± 0.05
85	0.33 f	14.4 ± 0.4	5.9 ± 0.5	0.67 f	19.4 ± 0.8	8.2 ± 1.0	10.9 ± 0.7	0.36 ± 0.06
86	0.33 f	14.6 ± 0.3	5.3 ± 0.4	0.67 f	19.4 ± 0.8	7.8 ± 0.8	10.4 ± 0.7	0.34 ± 0.06
87	0.33 f	14.7 ± 0.3	5.1 ± 0.4	0.67 f	19.3 ± 0.6	7.0 ± 0.6	9.6 ± 0.5	0.33 ± 0.05
88	0.33 f	14.8 ± 0.3	5.1 ± 0.4	0.67 f	19.4 ± 0.6	6.8 ± 0.6	9.5 ± 0.5	0.32 ± 0.05
89	0.33 f	14.8 ± 0.3	5.0 ± 0.4	0.67 f	19.4 ± 0.6	6.4 ± 0.6	9.2 ± 0.5	0.32 ± 0.05
90	0.33 f	14.7 ± 0.3	4.9 ± 0.4	0.67 f	19.2 ± 0.6	6.2 ± 0.6	8.9 ± 0.5	0.32 ± 0.05
91	0.33 f	14.7 ± 0.4	5.0 ± 0.4	0.67 f	19.0 ± 0.6	6.3 ± 0.8	8.8 ± 0.6	0.31 ± 0.05
92	0.33 f	14.8 ± 0.4	5.2 ± 0.4	0.67 f	19.0 ± 0.6	6.2 ± 0.8	8.7 ± 0.6	0.30 ± 0.06
93	$0.58^{+0.15}_{-0.35}$	$15.5^{+0.6}_{-1.2}$	5.52 ± 0.52	$0.32^{+0.33}_{-0.11}$	$18.8^{+0.9}_{-1.6}$	6.87 ± 2.36	7.63 ± 0.47	0.20 ± 0.05
94	$0.33^{+0.11}_{-0.20}$	$14.5^{+0.5}_{-0.8}$	4.68 ± 0.84	$0.57^{+0.22}_{-0.14}$	$17.8^{+0.6}_{-0.9}$	7.31 ± 1.17	8.60 ± 0.55	0.22 ± 0.03
95	0.22 ± 0.34	14.0 ± 1.7	4.78 ± 2.50	0.93 ± 0.36	17.9 ± 1.4	12.78 ± 2.37	12.08 ± 0.97	
96	$0.75^{+0.38}_{-2.22}$	$14.6^{+2.7}_{-4.7}$	6.70 ± 4.36	$0.88^{+2.22}_{-0.40}$	$17.9^{+2.4}_{-5.3}$	11.48 ± 2.70	9.79 ± 0.62	0.21 ± 0.27
97	$0.63^{+0.15}_{-1.30}$	$14.6^{+0.5}_{-3.9}$	7.20 ± 0.26	$0.47^{+0.7}_{-0.13}$	$18.3^{+0.7}_{-4.9}$	9.91 ± 1.04	9.90 ± 0.63	0.24 ± 0.03
98	$0.68^{+0.16}_{-0.27}$	$14.5^{+0.6}_{-1.0}$	7.16 ± 0.48	$0.35^{+0.25}_{-0.12}$	$18.8^{+0.8}_{-1.3}$	8.19 ± 1.61	10.06 ± 0.64	0.28 ± 0.02
99	1.0 ± 0.1	14.5 ± 0.5	9 ± 1					
100	1.00 f	15.00 f	9.00 f					
101		15.8 ± 0.5	8.5 ± 1					
102		15.4 ± 0.5	9.6 ± 1					
103	0.65	15.5 ± 0.5	6.0 ± 0.7					
104	0.65	15.5 ± 0.5	6.0 ± 0.7					
105	1.00 f	15.10	6.33					
106	0.63 f	15.05	5.94	0.37	15.91	6.45		-0.02 ± 0.02
107	1.00 f	15.12	5.79					
108	1.00 f	15.01	5.42					
109	0.65	15.5 ± 0.5	6.5 ± 0.7					
110	1.00 f	15.7 ± 0.5	11.4 ± 0.6					
111	1.00 f	15.7 ± 0.500	10.8 ± 0.600					
112	1.00 f	15.6 ± 0.500	11.4 ± 0.600					
113	0.90 ± 0.10	14.7 ± 0.2	7.3 ± 0.5					
114	0.90 ± 0.10	14.7 ± 0.2	7.3 ± 0.5					

No.	Ref.	Pro.	Tar.	CN	E_{lab}	E_{ex}	$\langle I_i \rangle$	T_i	l.d.	A/a	$\langle I_f \rangle$	FWHM	E_f	T_f	Code	gate
115	Bra95	^{48}Ti	^{62}Ni	^{110}Sn	223	92	40			8	45	24			C	f>9
116	Bra95	^{48}Ti	^{62}Ni	^{110}Sn	223	92	40	1.96		8	44	16		1.8 ± 0.13	C	γm
117	Bra95	^{48}Ti	^{62}Ni	^{110}Sn	223	92	40	1.74		8	54	14		1.6 ± 0.13	C	γm
118	Cha87a	^{16}O	^{94}Mo	^{110}Sn	140	110.2				9					C	γm
119	Bra89	^{40}Ar	^{70}Ge	^{110}Sn	400	230	40								C	er
120	Gaa86	^{20}Ne	^{91}Zr	^{111}Sn	100	66.4	27			8					C	γm
121	Gaa86	^{20}Ne	^{91}Zr	^{111}Sn	140	99.2	41			8					C	γm
122	Cha87a	^{19}F	^{93}Nb	^{112}Sn	80	62.2	21.3				16.3				C	f=12.5±2.0
123	Di601	^{19}F	^{93}Nb	^{112}Sn	75	62.2	20.9	2.23	P/R	9					C	γm
124	Cha87a	^{19}F	^{93}Nb	^{112}Sn	100	79.3	30.2			9	20.2				C	f=14.5±2.5
125	Cha87a	^{19}F	^{93}Nb	^{112}Sn	160	130.0	43.5			9	21.9				C	f=16.0±2.5
126	Sto89a	^4He	^{110}Cd	^{114}Sn	24	26	8		D	9					C	did
127	Sto89a	^4He	^{110}Cd	^{114}Sn	26				D	9					C	did
128	Bal94	^1H	^{115}In	^{116}Sn	50	58.8	8			8					C/P2	
129	Kel99	^{18}O	^{100}Mo	^{118}Sn	122	107			R					2.14	C	
130	Kel99	^{18}O	^{100}Mo	^{118}Sn	147	116			R					2.21	C	
131	Kel99	^{18}O	^{100}Mo	^{118}Sn	166	123			R					2.26	C	
132	Kel99	^{18}O	^{100}Mo	^{118}Sn	197	134			R					2.36	C	
133	Kel99	^{18}O	^{100}Mo	^{118}Sn	214	141			R					2.42	C	
134	Hec03	^{17}O	^{120}Sn	^{120}Sn	1360	25			R				9.7	1.0	C	is
135	Bau98	^4He	^{120}Sn	^{120}Sn	200	34.4			F					1.24	C	is
136	Bau98	^4He	^{120}Sn	^{120}Sn	200	44.4			F					1.54	C	is
137	Bau98	^4He	^{120}Sn	^{120}Sn	200	54.4			F					1.80	C	is
138	Bau98	^4He	^{120}Sn	^{120}Sn	200	64.4			F					2.02	C	is
139	Bau98	^4He	^{120}Sn	^{120}Sn	200	74.5			F					2.23	C	is
140	Bau98	^4He	^{120}Sn	^{120}Sn	200	84.5			F					2.43	C	is
141	Bau98	^4He	^{120}Sn	^{120}Sn	200	94.5			F					2.61	C	is
142	Bau98	^4He	^{120}Sn	^{120}Sn	200	104.5			F					2.79	C	is
143	Bau98	^4He	^{120}Sn	^{120}Sn	200	114.5			F					2.96	C	is
144	Bau98	^4He	^{120}Sn	^{120}Sn	200	124.6			F					3.12	C	is
145	End92	^{136}Xe	^{48}Ti	^{136}Xe	2516	200				8					C	dis
146	End92	^{136}Xe	^{48}Ti	^{136}Xe	2516	320				8					C	dis
147	End92	^{136}Xe	^{48}Ti	^{136}Xe	2516	420				8					C	dis
148	Gar83	^{12}C	^{115}In	^{127}Cs		48.4	16								C	
149	Gar83	^{18}O	^{109}Ag	^{127}Cs		48.4	8								C	
150	Voj89	^{12}C	^{112}Sn	^{124}Ba	90	71				9					C	
151	Voj89	^{12}C	^{112}Sn	^{124}Ba	127	105				9					C	
152	Voj89	^{12}C	^{124}Sn	^{136}Ba	90	83				9					C	
153	Voj89	^{12}C	^{124}Sn	^{136}Ba	127	116				9					C	
154	Di601	^{16}O	^{115}In	^{131}La	80	59.7	23.8	2.02	P/R	9					C	γm
155	Di601	^{19}F	^{115}In	^{134}Ce	75	58.1	16.9	1.98	P/R	9					C	γm
156	Kmi01	^{37}Cl	^{110}Pd	^{142}Eu	160	74	39								C	d γ t
157	Kmi01	^{37}Cl	^{110}Pd	^{142}Eu	170	81	43.0								C	d γ t
158	Kmi01	^{37}Cl	^{110}Pd	^{143}Eu	160	74	39								C	d γ t
159	Kmi01	^{37}Cl	^{110}Pd	^{143}Eu	170	81	43.0								C	d γ t
160	Kmi01	^{37}Cl	^{110}Pd	^{144}Eu	170	81	43.0								C	d γ t
161	Kmi01	^{37}Cl	^{110}Pd	^{144}Eu	160	74	39								C	d γ t
162	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	160	73.8	39		R		37			1.29	C	f=5–7
163	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	160	73.8	39		R		42			1.25	C	f=5–30
164	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	160	73.8	39		R		46			1.22	C	f=8–30
165	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	160	73.8	39		R		37			1.29	C	f=5–7
166	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	160	73.8	39		R		42			1.25	C	f=5–30
167	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	160	73.8	39		R		46			1.22	C	f=8–30
168	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	165	77.6	41		R		38			1.32	C	f=5–7
169	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	165	77.6	41		R		44			1.29	C	f=5–30
170	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	165	77.6	41		R		49			1.24	C	f=8–30
171	Kmi00	^{37}Cl	^{110}Pd	^{147}Eu	165	77.6	41		R		38			1.32	C	f=5–7

No.	S_1	E_1	Γ_1	S_2	E_2	Γ_2	FWHM	def.
115	1.00 f	14.9 ± 0.5	11.8 ± 0.6					
116	1.00 f	15.0 ± 0.500	11.7 ± 0.600					
117	1.00 f	14.7 ± 0.500	12.8 ± 0.600					
118	1.00 f	14.9 ± 0.3	9.8 ± 0.5					
119	1.00 f	16 ± 1	13^{+1}_{-2}					
120	1.0 ± 0.1	15.5	7.5					
121	1.0 ± 0.1	15.5 ± 0.5	11					
122	1.00 f	14.9 ± 0.3	6.7 ± 0.4					
123	1.00 f		6.5					
124	1.00 f	14.7 ± 0.3	7.9 ± 0.4					
125	1.00 f	14.6 ± 0.4	10.8 ± 0.6					
126	0.9 ± 0.1	15.2 ± 0.5	7 ± 1					
127	0.9 ± 0.1	15.2 ± 0.5	7 ± 1					
128	1.00 f	15.4 ± 0.7	6.2 ± 0.5					
129	1.2 ± 0.2	14.8 ± 0.5	8.3 ± 0.4					
130	1.1 ± 0.2	14.6 ± 0.5	8.7 ± 0.4					
131	1.3 ± 0.2	15.2 ± 0.5	10.0 ± 0.5					
132	1.2 ± 0.2	15.2 ± 0.5	10.7 ± 0.4					
133	1.2 ± 0.2	15.2 ± 0.5	10.6 ± 0.4					
134	1.1 ± 0.2	16.5 ± 0.7	4 ± 1					
135			5.5 ± 1.0					
136			7.5 ± 1.0					
137			7.5 ± 1.0					
138			8.5 ± 1.0					
139			9.0 ± 1.0					
140			9.5 ± 1.0					
141			10.0 ± 1.0					
142			10.0 ± 1.0					
143			12.0 ± 1.0					
144			11.5 ± 1.0					
145	1.00 f	15.3 f	9 ± 2					
146	1.00 f	15.3 f	10 ± 1.5					
147	1.00 f	15.3 f	10 ± 1					
148	1.3 f	14.25 ± 0.63	6.0 ± 0.63					
149	1.3 f	14.25 ± 0.63	6.0 ± 0.63					
150		15.0 ± 0.3	8.0 ± 0.4					
151		15.5 ± 0.3	8.5 ± 0.4					
152		14.9 ± 0.3	7.8 ± 0.4					
153		15.2 ± 0.3	8.3 ± 0.4					
154	1.00 f		6.5					
155	1.00 f		6.5					
156	0.9 ± 0.1	14.3 f	8.8 ± 0.8					
157	1.0 ± 0.1	14.3 f	9.8 ± 1.0					
158	1.0 ± 0.1	14.3 f	9.3 ± 0.4					
159	1.0 ± 0.1	14.3 f	9.8 ± 0.5					
160	0.9 ± 0.2	14.3 f	12.0 ± 0.8					
161	1.0 ± 0.2	14.3 f	8.8 ± 0.7					
162	0.83 ± 0.02	14.3 ± 0.1	8.0 ± 0.2					
163	0.88 ± 0.02	14.3 ± 0.1	8.4 ± 0.2					
164	0.86 ± 0.03	14.2 ± 0.1	8.7 ± 0.4					
165	0.36 ± 0.02	13.1 ± 0.1	5.8 ± 0.1	0.46 ± 0.03	16.4 ± 0.1	7.7 ± 0.3		$\pm 0.24 \pm 0.01$
166	0.39 ± 0.02	13.0 ± 0.1	5.9 ± 0.1	0.45 ± 0.03	16.6 ± 0.1	7.7 ± 0.2		$\pm 0.27 \pm 0.01$
167	0.42 ± 0.02	12.9 ± 0.1	5.9 ± 0.1	0.38 ± 0.04	16.8 ± 0.2	7.5 ± 0.5		$\pm 0.29 \pm 0.02$
168	1.01 ± 0.03	14.2 ± 0.1	8.0 ± 0.2					
169	1.00 ± 0.02	14.3 ± 0.1	8.4 ± 0.2					
170	1.10 ± 0.03	14.2 ± 0.1	8.6 ± 0.3					
171	0.49 ± 0.02	13.0 ± 0.1	5.9 ± 0.1	0.51 ± 0.03	16.5 ± 0.1	7.3 ± 0.3		$\pm 0.26 \pm 0.01$

No.	Ref.	Pro.	Tar.	CN	E_{lab}	E_{ex}	$\langle I_i \rangle$	T_i	l.d.	A/a	$\langle I_f \rangle$	FWHM	E_f	T_f	Code	gate
172	Kmi00	³⁷ Cl	¹¹⁰ Pd	¹⁴⁷ Eu	165	77.6	41		R		44			1.29	C	f=5–30
173	Kmi00	³⁷ Cl	¹¹⁰ Pd	¹⁴⁷ Eu	165	77.6	41		R		49			1.24	C	f=8–30
174	Kmi00	³⁷ Cl	¹¹⁰ Pd	¹⁴⁷ Eu	170	81.4	43.0		R		39			1.36	C	f=5–7
175	Kmi00	³⁷ Cl	¹¹⁰ Pd	¹⁴⁷ Eu	170	81.4	43.0		R		45			1.31	C	f=5–30
176	Kmi00	³⁷ Cl	¹¹⁰ Pd	¹⁴⁷ Eu	170	81.4	43.0		R		50			1.28	C	f=8–30
177	Kmi00	³⁷ Cl	¹¹⁰ Pd	¹⁴⁷ Eu	170	81.4	43.0		R		39			1.36	C	f=5–7
178	Kmi00	³⁷ Cl	¹¹⁰ Pd	¹⁴⁷ Eu	170	81.4	43.0		R		45			1.31	C	f=5–30
179	Kmi00	³⁷ Cl	¹¹⁰ Pd	¹⁴⁷ Eu	170	81.4	43.0		R		50			1.28	C	f=8–30
180	Haa83	²⁸ Si	¹²⁰ Sn	¹⁴⁴ Gd	145											did & f $\geq$ 3
181	Haa83	²⁸ Si	¹²⁰ Sn	¹⁴⁴ Gd	145											did & f $\geq$ 3
182	Fli96	³⁰ Si	¹¹⁶ Sn	¹⁴⁶ Gd	156	82					22				C	lcp & f=4–8
183	Fli96	³⁰ Si	¹¹⁶ Sn	¹⁴⁶ Gd	156	82					36				C	lcp & f=9–12
184	Fli96	³⁰ Si	¹¹⁶ Sn	¹⁴⁶ Gd	156	82					58				C	lcp & f=13–20
185	Fli96	⁷⁶ Ge	⁷⁰ Ge	¹⁴⁶ Gd	319	82					22				C	lcp & f=4–8
186	Fli96	⁷⁶ Ge	⁷⁰ Ge	¹⁴⁶ Gd	319	82					36				C	lcp & f=9–12
187	Fli96	⁷⁶ Ge	⁷⁰ Ge	¹⁴⁶ Gd	319	82					58				C	lcp & f=13–20
188	Fli91	³⁰ Si	¹²⁴ Sn	¹⁵⁴ Gd	150	80	47			9	27				C/P	f=10–16
189	Fli91	³⁰ Si	¹²⁴ Sn	¹⁵⁴ Gd	150	80	47			9	35				C/P	f=17–20
190	Fli91	³⁰ Si	¹²⁴ Sn	¹⁵⁴ Gd	150	80	47			9	43				C/P	f=21–25
191	Fli91	³⁰ Si	¹²⁴ Sn	¹⁵⁴ Gd	150	80	47			9	55				C/P	f=26–36
192	Noo92	⁴⁰ Ar	¹¹⁴ Cd	¹⁵¹ Dy	173	69	50			8	40		1.60		C	did & f $\geq$ 2
193	Noo92	⁴⁰ Ar	¹¹⁴ Cd	¹⁵¹ Dy	173	69	50			8	40		1.60		C	did & f $\geq$ 2
194	Noo92	⁴⁰ Ar	¹¹⁴ Cd	¹⁵¹ Dy	173	69	50			8	40		1.60		C	did & f $\geq$ 2
195	Noo92	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	173	69	50			8	40				C	
196	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	32				C	γm & $\gamma \Sigma E$
197	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	41				C	γm & $\gamma \Sigma E$
198	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	50				C	γm & $\gamma \Sigma E$
199	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			8	32				C	γm & $\gamma \Sigma E$
200	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			8	41				C	γm & $\gamma \Sigma E$
201	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			8	50				C	γm & $\gamma \Sigma E$
202	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			8	32				C	γm & $\gamma \Sigma E$
203	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			8	41				C	γm & $\gamma \Sigma E$
204	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			8	50				C	γm & $\gamma \Sigma E$
205	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	32				C	γm & $\gamma \Sigma E$
206	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	41				C	γm & $\gamma \Sigma E$
207	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	50				C	γm & $\gamma \Sigma E$
208	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	32				C	γm & $\gamma \Sigma E$
209	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	41				C	γm & $\gamma \Sigma E$
210	Noo94	⁴⁰ Ar	¹¹⁴ Cd	¹⁵⁴ Dy	167	69	50			9	50				C	γm & $\gamma \Sigma E$
211	Bru88	⁴⁸ Ca	¹⁰⁸ Pd	¹⁵⁶ Dy	206	75	40				18	10			C	γm
212	Bru88	⁴⁸ Ca	¹⁰⁸ Pd	¹⁵⁶ Dy	206	75	40				28	10			C	γm
213	Bru88	⁴⁸ Ca	¹⁰⁸ Pd	¹⁵⁶ Dy	206	75	40				40	10			C	γm
214	Bru88	⁴⁸ Ca	¹⁰⁸ Pd	¹⁵⁶ Dy	206	75	40				18	10			C	γm
215	Bru88	⁴⁸ Ca	¹⁰⁸ Pd	¹⁵⁶ Dy	206	75	40				28	10			C	γm
216	Bru88	⁴⁸ Ca	¹⁰⁸ Pd	¹⁵⁶ Dy	206	75	40				40	10			C	γm
217	Sto89b	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	205	90	41			8	32				C	γm & $\gamma \Sigma E$
218	Sto89b	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	205	90	41			8	46				C	γm & $\gamma \Sigma E$
219	Sto89b	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	205	90	41			8	62				C	γm & $\gamma \Sigma E$
220	Sch93	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	196	92.5	60	1.89		8	32	20		1.64	C	γm & $\gamma \Sigma E$
221	Sch95	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	200	92.5	75			8	32	20		1.8 $\pm$ 0.2	C	st
222	Sch93	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	196	92.5	60	1.78		8	46	20		1.54	C	γm & $\gamma \Sigma E$
223	Sch95	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	200	92.5	75			8	46	20		1.7 $\pm$ 0.2	C	st
224	Sch93	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	196	92.5	60	1.60		8	62	20		1.35	C	γm & $\gamma \Sigma E$
225	Hof98	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	216.0	104.3	61			10	82				C	ff
226	Hof98	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	216.0	104.3	61			10	82				C	ff
227	Hof98	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	216.0	104.3	61			10	82				C	ff
228	Hof98	⁴⁰ Ar	¹¹⁶ Cd	¹⁵⁶ Dy	216.0	104.3	61			10	82				C	ff

No.	S_1	E_1	Γ_1	S_2	E_2	Γ_2	FWHM	def.
172	0.42 ± 0.02	13.0 ± 0.1	5.8 ± 0.1	0.43 ± 0.02	16.7 ± 0.1	7.4 ± 0.2		$\pm 0.27 \pm 0.01$
173	0.54 ± 0.02	12.9 ± 0.1	5.8 ± 0.1	0.48 ± 0.03	17.0 ± 0.1	7.1 ± 0.4		$\pm 0.3 \pm 0.01$
174	0.91 ± 0.03	14.2 ± 0.1	8.8 ± 0.4					
175	0.98 ± 0.03	14.1 ± 0.1	8.9 ± 0.3					
176	1.01 ± 0.04	14.0 ± 0.1	8.8 ± 0.3					
177	0.43 ± 0.02	12.9 ± 0.1	5.9 ± 0.1	0.41 ± 0.03	16.8 ± 0.1	7.3 ± 0.4		$\pm 0.29 \pm 0.01$
178	0.48 ± 0.02	12.9 ± 0.1	5.9 ± 0.1	0.40 ± 0.03	16.8 ± 0.1	7.0 ± 0.3		$\pm 0.29 \pm 0.01$
179	0.52 ± 0.02	12.9 ± 0.1	5.9 ± 0.1	0.40 ± 0.04	17.0 ± 0.2	7.2 ± 0.5		$\pm 0.3 \pm 0.02$
180		11.6	6.7	$S_1/2$	14.9	6.7		
181		12.2	8.9					
182		13.8	8.4					
183		13.8	8.4					
184		13.8	8.4					
185		13.8	8.4					
186		13.8	8.4					
187		13.8	8.4					
188	$0.30 f$	13.5 ± 0.5	7.5 ± 1.0	0.70 ± 0.1	15.3 ± 0.5	$9.6 f$		0.08 ± 0.08
189	$0.30 f$	13.5 ± 0.5	7.5 ± 1.0	0.70 ± 0.1	15.3 ± 0.5	$9.6 f$		0.18 ± 0.08
190	$0.30 f$	12.5 ± 0.5	5.5 ± 1.0	0.70 ± 0.1	15.5 ± 0.5	$8.5 f$		0.23 ± 0.08
191	$0.40 f$	12.0 ± 0.5	5.5 ± 1.0	0.60 ± 0.1	16.0 ± 0.5	$9.8 f$		0.35 ± 0.08
192	$1.00 f$	14.0 ± 0.5	8.4 ± 1.2					
193	$0.38 f$	12.0 ± 0.5	$3.5 \pm 0.8^*$	$0.62 f$	17.6 ± 0.7	$7.4 \pm 1.7^*$		0.40 ± 0.005
194	$0.75 f$	13.2 ± 0.5	$5.2 \pm 1.0^*$	$0.25 f$	21.3 ± 0.7	$13.6 \pm 2.4^*$		-0.50 ± 0.005
195	$1.00 f$	15.5 ± 0.5	10.6 ± 1.0					
196	$1.00 f$	15.0 ± 0.3	9.5 ± 1.2					
197	$1.00 f$	15.1 ± 0.3	10.3 ± 1.2					
198	$1.00 f$	15.2 ± 0.3	14.6 ± 1.3					
199	$1.00 f$	14.6 ± 0.3	7.6					
200	$1.00 f$	14.7 ± 0.3	8.4					
201	$1.00 f$	15.1 ± 0.3	11.8					
202	0.47	14.4 ± 0.3	7.4	$0.53 f$	15.5 ± 0.7	8.5		± 0.11
203	0.38	14.9 ± 0.3	8.8	$0.62 f$	15.3 ± 0.7	9.3		0.13
204	0.67	15.0 ± 0.3	11.6	$0.33 f$	15.1 ± 0.7	11.7		-0.18
205	$0.35 f$	14.0 ± 0.3	$8.0 \pm 1.0^*$	$0.65 f$	15.7 ± 0.7	$10.1 \pm 1.5^*$		0.11
206	$0.35 f$	13.5 ± 0.3	$7.7 \pm 1.0^*$	$0.65 f$	16.5 ± 0.7	$11.4 \pm 1.7^*$		0.13
207	$0.35 f$	13.6 ± 0.3	$11.3 \pm 1.1^*$	$0.65 f$	16.4 ± 0.7	$16.4 \pm 1.9^*$		0.18
208	$0.65 f$	14.5 ± 0.3	$8.4 \pm 1.1^*$	$0.35 f$	16.5 ± 0.7	$10.9 \pm 1.6^*$		-0.11
209	$0.65 f$	14.4 ± 0.3	$9.1 \pm 1.1^*$	$0.35 f$	17.0 ± 0.7	$12.7 \pm 1.8^*$		-0.13
210	$0.65 f$	14.1 ± 0.3	$11.3 \pm 1.1^*$	$0.35 f$	20.0 ± 0.7	$22.8 \pm 2.6^*$		-0.18
211	$1.00 f$	13.9 ± 0.5	10.9					
212	$1.00 f$	13.4 ± 0.5	10.3					
213	$1.00 f$	13.4 ± 0.5	9.2					
214	0.28 ± 0.05	10.8	10.7	0.72 ± 0.05	14.6	8.7		0.36 ± 0.05
215	0.60 ± 0.05	12.3	10.8	0.40 ± 0.05	14.4	7.3		-0.18 ± 0.05
216	0.65 ± 0.05	12.7	8.3	0.35 ± 0.05	15.0	9.2		-0.19 ± 0.05
217	0.28	13.8	8.6 ± 0.5	$0.72 f$	15.8	$11.3 f$		0.16 ± 0.04
218	0.68	13.9	8.5 ± 0.5	$0.32 f$	15.5	$10.6 f$		-0.13 ± 0.04
219	0.67	13.5	8.5 ± 0.5	$0.33 f$	17.1	$13.6 f$		-0.28 ± 0.04
220	$0.70 \pm 0.0045^*$	14.4 ± 1.0	$9.7 \pm 1.4^*$	$0.30 \pm 0.0045^*$	17.1 ± 0.5	$13.8 \pm 1.0^*$		$^{(+)} 0.20 \pm 0.06$
221	$0.34 \pm 0.05^*$	11.4 ± 1.6	$7.1 \pm 2.2^*$	$0.66 \pm 0.05^*$	17.2 ± 1.4	$16.3 \pm 3.6^*$		$\pm 0.51 \pm 0.29$
222	$0.45 \pm 0.01^*$	12.7 ± 1.2	$6.1 \pm 1.6^*$	$0.55 \pm 0.01^*$	16.9 ± 1.1	$10.9 \pm 2.4^*$		$\pm 0.34 \pm 0.06$
223	$0.50 \pm 0.09^*$	12.2 ± 1.4	$7.0 \pm 2.0^*$	$0.50 \pm 0.09^*$	16.8 ± 1.2	$13.2 \pm 2.9^*$		$\mp 0.35 \pm 0.14$
224	$0.50 \pm 0.09^*$	12.2 ± 1.1	$6.3 \pm 1.3^*$	$0.50 \pm 0.09^*$	16.8 ± 0.9	$11.8 \pm 1.7^*$		$\pm 0.35 \pm 0.06$
225	0.26 ± 0.01	10.10 ± 0.04	$3.5 \pm 1.8^*$	0.99 ± 0.02	15.26 ± 0.05	$7.9 \pm 4.2^*$		0.499 ± 0.003
226	0.34 ± 0.32	10.38 ± 0.94	$3.0 \pm 2.6^*$	0.96 ± 0.32	15.54 ± 1.41	$6.8 \pm 5.9^*$		0.488 ± 0.007
227	0.26 ± 0.01	10.10 ± 0.03	$3.5 \pm 1.2^*$	0.99 ± 0.01	15.26 ± 0.04	$7.9 \pm 2.8^*$		0.499 ± 0.003
228	0.20 ± 0.06	10.76 ± 0.04	$5.7 \pm 0.1^*$	0.80 ± 0.24	16.24 ± 0.05	$13.1 \pm 0.2^*$		0.498 ± 0.003

No.	Ref.	Pro.	Tar.	CN	E_{lab}	E_{ex}	$\langle I_i \rangle$	T_i	l.d.	A/a	$\langle I_f \rangle$	FWHM	E_f	T_f	Code	gate
229	Hof98	^{40}Ar	^{116}Cd	^{156}Dy	242.4	124.0	70			10	89				C	ff
230	Hof98	^{40}Ar	^{116}Cd	^{156}Dy	242.4	124.0	70			10	89				C	ff
231	Hof98	^{40}Ar	^{116}Cd	^{156}Dy	242.4	124.0	70			10	89				C	ff
232	Hof98	^{40}Ar	^{116}Cd	^{156}Dy	242.4	124.0	70			10	89				C	ff
233	Vie96	^{64}Ni	^{92}Zr	^{154}Er	241					8	17.5	25			C	d γ t
234	Vie97	^{64}Ni	^{92}Zr	^{154}Er	241		31								C	d γ t
235	Vie97	^{64}Ni	^{92}Zr	^{155}Er	241		31								C	d γ t
236	Vie96	^{64}Ni	^{92}Zr	^{156}Er	241					8	25	18			C	f=11–15
237	Vie96	^{64}Ni	^{92}Zr	^{156}Er	241					8	35.5	21			C	f=16–20
238	Vie96	^{64}Ni	^{92}Zr	^{156}Er	241					8	41.5	13			C	f=21–30
239	Gaa88	^{16}O	^{142}Nd	^{158}Er		47				8	12					$\gamma\Sigma\text{E}$
240	Gaa88	^{16}O	^{142}Nd	^{158}Er		47				8	20					$\gamma\Sigma\text{E}$
241	Gaa88	^{16}O	^{142}Nd	^{158}Er		47				8	26					$\gamma\Sigma\text{E}$
242	Gos85	^{12}C	^{148}Sm	^{160}Er	63	43.2	15			8	12.5	25	30	1	C	
243	Cha88	^{19}F	^{141}Pr	^{160}Er	94.1	59.2				9	25			1.3	C	γm
244	Cha88	^{19}F	^{141}Pr	^{160}Er	94.1	59.2				9	25			1.3	C	γm
245	Cha88	^{19}F	^{141}Pr	^{160}Er	94.1	59.2				9	25			1.3	C	γm
246	Cha88	^{19}F	^{141}Pr	^{160}Er	96.5	61.2				9	25			1.3	C	γm
247	Cha88	^{19}F	^{141}Pr	^{160}Er	96.5	61.2				9	25			1.3	C	γm
248	Cha88	^{19}F	^{141}Pr	^{160}Er	96.5	61.2				9	25			1.3	C	γm
249	Nan99	^{40}Ar	^{124}Sn	^{160}Er	160	62	31			7.5	44		13.8	1	C	er
250	Cha88	^{19}F	^{141}Pr	^{160}Er	111.1	74.3				9	34			1.4	C	γm
251	Cha88	^{19}F	^{141}Pr	^{160}Er	111.1	74.3				9	34			1.4	C	γm
252	Cha88	^{19}F	^{141}Pr	^{160}Er	111.1	74.3				9	34			1.4	C	γm
253	Cha88	^{19}F	^{141}Pr	^{160}Er	129.0	90.3				9	42			1.5	C	γm
254	Cha88	^{19}F	^{141}Pr	^{160}Er	129.0	90.3				9	42			1.5	C	γm
255	Cha88	^{19}F	^{141}Pr	^{160}Er	129.0	90.3				9	42			1.5	C	γm
256	Thi88	^{34}S	^{130}Te	^{164}Er	156	72	40			9	31				C	f=15–19
257	Thi88	^{34}S	^{130}Te	^{164}Er	156	72	40			9	39				C	f=20–24
258	Thi88	^{34}S	^{130}Te	^{164}Er	156	72	40			9	48				C	f=25–35
259	Bra96	^{17}O	^{148}Nd	^{165}Er	70	45	16							1.3	C	
260	Gos85	^{12}C	^{154}Sm	^{166}Er	63	49.2	15			8	12.5	25	30	1	C	
261	Bra96	^{18}O	^{148}Nd	^{166}Er	78	51					23			1.3	C	
262	Gaa88	^{16}O	^{150}Nd	^{166}Er		60				8	12					$\gamma\Sigma\text{E}$
263	Gaa88	^{16}O	^{150}Nd	^{166}Er		60				8	20					$\gamma\Sigma\text{E}$
264	Gaa88	^{16}O	^{150}Nd	^{166}Er		60				8	26					$\gamma\Sigma\text{E}$
265	Gaa84	^{16}O	^{150}Nd	^{166}Er	84	61.5				7.5					C	γc
266	Tho93	^{16}O	^{148}Sm	^{164}Yb	82	49				10	20	20			C	f=5–10
267	Tho92	^{16}O	^{159}Tb	^{175}Ta	160	123.4				10					C	er
268	Tho92	^{16}O	^{159}Tb	^{175}Ta	160	123.4				10	70	20		1.8	C	ff
269	Mat95	^{28}Si	^{148}Nd	^{176}W	147	70									C	f $\geq$ 9
270	Mat95	^{28}Si	^{148}Nd	^{176}W	147					8	36	16		1.49	C	γm
271	Mat95	^{28}Si	^{148}Nd	^{176}W	147					8	42	15		1.45	C	γm
272	Mat95	^{28}Si	^{148}Nd	^{176}W	147					8	47	15		1.41	C	γm
273	Mat95	^{28}Si	^{148}Nd	^{176}W	147					8	55	19		1.35	C	γm
274	Cam03	^{90}Zr	^{89}Y	^{179}Au	346	20	11			8				0.7		$\alpha\text{d of er \& } \gamma\text{m}$
275	Ata90	^{48}Ca	^{142}Nd	^{188}Hg	205	52	19			9					C	er & γm
276	Ata90	^{48}Ca	^{142}Nd	^{188}Hg	205	52	19			9					C	er & γm
277	Ata90	^{48}Ca	^{142}Nd	^{190}Hg	205	52	19			9					C	any er & γm
278	Ata90	^{48}Ca	^{142}Nd	^{190}Hg	205	52	19			9					C	any er & γm
279	Cam99	^{30}Si	^{164}Dy	^{191}Hg	142	60	27			8	24			1.3		f $\geq$ 2–5 & d γ t
280	Cam99	^{30}Si	^{164}Dy	^{191}Hg	142	60	27			8	36			1.3		f>7 & d γ t
281	Cam99	^{30}Si	^{164}Dy	^{194}Hg	142	60	27			8	27			1.3		f $\geq$ 2 & any d γ t
282	Cha96	^{16}O	^{181}Ta	^{197}Tl	83.6	50.4	13	I							C	γc
283	Cha96	^{16}O	^{181}Ta	^{197}Tl	88.7	55.2	20	I							C	γc
284	Cha96	^{16}O	^{181}Ta	^{197}Tl	94.1	60.1	24	I			15	8			C	f=3–4
285	Cha96	^{16}O	^{181}Ta	^{197}Tl	94.1	60.1	24	I			23.5	7			C	f=5–6

No.	S_1	E_1	Γ_1	S_2	E_2	Γ_2	FWHM	def.
229	0.21 ± 0.01	11.20 ± 0.14	$4.6 \pm 1.5^*$	0.84 ± 0.02	16.21 ± 0.06	$9.7 \pm 3.2^*$		0.446 ± 0.018
230	0.24 ± 0.03	11.03 ± 0.14	$3.8 \pm 0.5^*$	0.91 ± 0.09	15.97 ± 0.13	$7.9 \pm 1.0^*$		0.446 ± 0.015
231	0.21 ± 0.04	11.40 ± 0.06	$5.0 \pm 0.2^*$	0.86 ± 0.14	16.14 ± 0.02	$9.9 \pm 0.4^*$		0.419 ± 0.008
232	0.21 ± 0.01	11.94 ± 0.10	$6.7 \pm 0.2^*$	0.80 ± 0.01	16.01 ± 0.05	$12.0 \pm 0.2^*$		0.352 ± 0.012
233	0.90 ± 0.30	15.1 ± 0.6	9.0 ± 1.0					
234	1 f	15 f	7 f					
235	1 f	15 f	7 f					
236	1.20 ± 0.10	15.0 ± 0.5	6.0 ± 0.2					
237	0.95 ± 0.20	14.8 ± 0.6	6.8 ± 0.9					
238	1.20 ± 0.40	14.8 ± 0.7	8.1 ± 2.0					
239	$0.58 f$	13.90	5.68	0.42	16.07	5.96		-0.16 ± 0.02
240	$0.68 f$	13.39	4.69	0.32	16.59	3.62		-0.24 ± 0.02
241	$0.70 f$	13.38	5.59	0.30	16.76	5.24		-0.26 ± 0.02
242	0.39 ± 0.21	12.21 ± 0.09	3.13 ± 0.21	0.64 ± 0.22	15.17 ± 0.18	4.81 ± 0.51		0.25
243	$0.33 f$	12.4 ± 0.2	4.9 ± 0.2	$0.67 f$	15.7 ± 0.3	6.9 ± 0.3		0.28 ± 0.03
244	$0.67 f$	13.1 ± 0.2	6.0 ± 0.2	$0.33 f$	16.6 ± 0.3	5.0 ± 0.3		-0.27 ± 0.03
245	$0.33 f$	12.3 ± 0.2	5.3 ± 0.3	triaxial all S + Γ eq.	14.1 ± 0.2	$E_3 = 16.9 \pm 0.3$	$\gamma = 32(4)^\circ$	0.29 ± 0.04
246	$0.33 f$	12.4 ± 0.2	4.8 ± 0.2	$0.67 f$	15.5 ± 0.3	7.5 ± 0.3		0.27 ± 0.03
247	$0.67 f$	12.9 ± 0.2	5.8 ± 0.2	$0.33 f$	16.3 ± 0.3	5.3 ± 0.3		-0.26 ± 0.03
248	$0.33 f$	12.2 ± 0.2	5.3 ± 0.3	triaxial all S + Γ eq.	13.6 ± 0.2	$E_3 = 16.6 \pm 0.3$	$\gamma = 37(4)^\circ$	0.28 ± 0.04
249	$0.4 \pm 0.05^*$	11.9 ± 0.3	3.6 ± 0.4	$0.6 \pm 0.07^*$	15.1 ± 0.3	6.4 ± 0.8		0.27 ± 0.02
250	$0.33 f$	12.4 ± 0.2	5.3 ± 0.3	$0.67 f$	15.3 ± 0.3	7.9 ± 0.4		0.25 ± 0.03
251	$0.67 f$	12.8 ± 0.2	6.0 ± 0.3	$0.33 f$	16.1 ± 0.3	5.8 ± 0.4		-0.26 ± 0.03
252	$0.33 f$	12.3 ± 0.2	5.8 ± 0.4	triaxial all S + Γ eq.	13.3 ± 0.3	$E_3 = 16.2 \pm 0.3$	$\gamma = 42(6)^\circ$	0.25 ± 0.04
253	$0.33 f$	12.4 ± 0.2	6.1 ± 0.3	$0.67 f$	15.2 ± 0.3	9.3 ± 0.4		0.24 ± 0.03
254	$0.67 f$	12.8 ± 0.2	6.9 ± 0.3	$0.33 f$	16.3 ± 0.3	7.3 ± 0.4		-0.27 ± 0.03
255	$0.33 f$	12.6 ± 0.3	7.0 ± 0.4	triaxial all S + Γ eq.	12.9 ± 0.3	$E_3 = 16.2 \pm 0.4$	$\gamma = 55(8)^\circ$	0.24 ± 0.04
256	$0.67 f$	13.4	4.5	$0.33 f$	14.6	4.5 f		-0.1
257	$0.67 f$	11.8	4.0	$0.33 f$	15.4	4.5 f		-0.3
258	$0.67 f$	11.6	4.0	$0.33 f$	15.2	4.5 f		-0.3
259	$0.60 f$	12.5 f	4.3 f	0.4 f	15.2 f	4.0 f		
260	0.43 ± 0.21	12.15 ± 0.09	3.69 ± 0.23	0.74 ± 0.23	15.77 ± 0.17	5.75 ± 0.71		0.30
261	$0.60 f$	12.5 f	4.6 f	0.4 f	15.2 f	4.2 f		
262	$0.32 f$	12.33	5.91	0.68	15.12	6.00		0.23 ± 0.02
263	$0.54 f$	12.79	5.87	0.46	15.32	6.22		$\pm 0.2 \pm 0.02$
264	$0.60 f$	12.93	5.99	0.40	16.65	5.07		$\pm 0.22 \pm 0.02$
265	0.60	12.6	5.2 f	0.40	16.4	5.2 f		$\pm 0.27 \pm 0.07$
266	0.33	12.2 ± 0.2	4.9 ± 0.5	0.67	15.7 ± 0.5	6.9 ± 1.0		
267		12.4	8		15.3	10.5		
268		9.5 ± 0.5	5 ± 1.0	2S ₁ f	15.0 ± 1.0	7.0 ± 2.0		0.55 ± 0.11
269		14.0	8.8					
270	1.00 f	14.2 ± 0.500	8.9 ± 0.600					
271	1.00 f	14.1 ± 0.500	8.4 ± 0.600					
272	1.00 f	13.6 ± 0.500	8.6 ± 0.600					
273	1.00 f	13.5 ± 0.500	8.9 ± 0.800					
274	1.00 f	14.2 f	$\Gamma_0 = 5 \pm 0.35$	spherical Hill-Wheeler		$\Gamma_0(E/E_0)^{1.6}$		0.10 ± 0.10
275	1.00 f	14.6 ± 0.5	7.0 ± 0.7					
276	0.34 ± 0.06	12.3 ± 0.5	5.0 ± 0.8	0.66 ± 0.06	15.5 ± 0.5	5.0 ± 0.8		0.28 ± 0.05
277	1.00 f	14.8 ± 0.1	10.4 ± 0.6					
278	0.44 ± 0.05	12.8 ± 0.1	7.6 ± 0.5	0.56 ± 0.05	16.6 ± 0.2	8.3 ± 0.4		0.31 ± 0.02
279	1.00 f	14	6.5					0.3
280	1.00 f	14	5.5					0.3
281	1.00 f	14.0	6.5					0.3
282	$0.33 f$	11.4 ± 0.2	4.0 ± 0.5	$0.67 f$	14.2 ± 0.7	6.2 ± 0.7		0.27 ± 0.04
283	$0.33 f$	11.5 ± 0.2	5.0 ± 0.5	$0.67 f$	14.4 ± 0.7	7.0 ± 0.7		$0.27^{+0.04}_{-0.02}$
284	$0.33 f$	11.7 ± 0.2	3.9 ± 0.5	$0.67 f$	15.4 ± 0.7	5.4 ± 0.7		0.26 ± 0.01
285	$0.33 f$	11.6 ± 0.2	3.5 ± 0.5	$0.67 f$	15.2 ± 0.7	5.2 ± 0.7		$0.26^{+0.03}_{-0.02}$

No.	Ref.	Pro.	Tar.	CN	E_{lab}	E_{ex}	$\langle I_i \rangle$	T_i	l.d.	A/a	$\langle I_f \rangle$	FWHM	E_f	T_f	Code	gate
286	Cha96	^{16}O	^{181}Ta	^{197}Tl	94.1	60.1	24		I		28				C	f=7–14
287	Cha96	^{16}O	^{181}Ta	^{197}Tl	106.5	71.6	33		I						C	γc
288	Tho88	^{19}F	^{181}Ta	^{200}Pb	93	57.6	14			8	7	14		0.9	C	
289										8	18.5	7		0.9	C	
290	Gel00	^{19}F	^{181}Ta	^{200}Pb	101	65	21			8					C	$f \geq 6$
291	Cha87b	^{16}O	natW	Pb	100	65.7	25			8				1.1	C	γm
292	Cha87b	^{19}F	^{181}Ta	^{200}Pb	105	68.5				8				1.1	C	γm
293	But90	^{19}F	^{181}Ta	^{200}Pb	105	69.5	34		P	8				1.2	C	ff
294	But90	^{19}F	^{181}Ta	^{200}Pb	105	69.5	25		P	8				1.1	C	γm
295	Sch96	^{19}F	^{181}Ta	^{200}Pb	95	86									C	γm & ipc
296	Cha87b	^{19}F	^{181}Ta	^{200}Pb	126	87.8	38			8				1.3	C	γm
297	But90	^{19}F	^{181}Ta	^{200}Pb	141	102.4	50		P	8				1.5	C	γm or ff
298	But90	^{19}F	^{181}Ta	^{200}Pb	141	102.4	50		P	8				1.5	C	γm or ff
299	Cha87b	^{16}O	natW	Pb	140	102.5	44			8				1.5	C	γm
300	Cha87b	^{16}O	natW	Pb	140	102.5	44			8				1.5	C	γm
301	Dió00a	^{19}F	^{181}Ta	^{200}Pb	161	121	49	2.2	R	9					C	ff
302	Dió00a	^{19}F	^{181}Ta	^{200}Pb	181	139	56	2.4	R	9					C	ff
303	Bau98	^4He	^{208}Pb	^{208}Pb	160.0	44.4			R					1.34	C	is
304	Bau98	^4He	^{208}Pb	^{208}Pb	160.0	54.4			R					1.49	C	is
305	Bau98	^4He	^{208}Pb	^{208}Pb	160.0	64.4			R					1.62	C	is
306	Bau98	^4He	^{208}Pb	^{208}Pb	160.0	74.4			R					1.74	C	is
307	Bau98	^4He	^{208}Pb	^{208}Pb	160.0	84.5			R					1.85	C	is
308	Bau98	^4He	^{208}Pb	^{208}Pb	160.0	94.6			R					1.95	C	is
309	Bau98	^4He	^{208}Pb	^{208}Pb	160.0	104.6			R					2.05	C	is
310	Bra05	^{18}O	^{198}Pt	^{212}Rn	96		38								C	did
311	Bra05	^{18}O	^{198}Pt	^{216}Rn	96		23								C	f=5–8
312	Bra05	^{18}O	^{198}Pt	^{216}Rn	96		29								C	f=9–30
313	Dió92	^{32}S	natW	Th	185	72	35	1.7		8.8					C	both ff
314	Dió92	^{32}S	natW	Th	185	72	35	1.7		8.8					C	both ff
315	Dió92	^{32}S	natW	Th	215	97	49	2.0		8.8					C	both ff
316	Dió92	^{32}S	natW	Th	215	97	49	2.0		8.8					C	both ff
317	Dió92	^{32}S	natW	Th	230	110	59	2.1		8.8					C	both ff
318	Dió92	^{32}S	natW	Th	230	110	59	2.1		8.8					C	both ff
319	Tho87	^{16}O	^{208}Pb	^{224}Th	100	44.0	25		P	8.8				0.9	C	
320	Dió00b	^{16}O	^{208}Pb	^{224}Th	100	46.4	25.0	1.37	R	9					C	ff
321	Tho87	^{16}O	^{208}Pb	^{224}Th	120	64.0	37		P	9.0				1.2	C	ff
322	Dió00b	^{16}O	^{208}Pb	^{224}Th	120	64.9	36.4	1.61	R	9					C	ff
323	Dió92	^{16}O	^{208}Pb	^{224}Th	140	82	43	1.8		8.8					C	both ff
324	Dió92	^{16}O	^{208}Pb	^{224}Th	140	82	43	1.8		8.8					C	both ff
325	Tho87	^{16}O	^{208}Pb	^{224}Th	140	82.0	43		P	8.8				1.4	C	
326	Dió00b	^{16}O	^{208}Pb	^{224}Th	140	83.5	44.5	1.83	R	9					C	ff
327	Dió00b	^{16}O	^{208}Pb	^{224}Th	160	102.1	50.3	2.03	R	9					C	ff
328	Dió00b	^{16}O	^{208}Pb	^{224}Th	177	117.9	54.1	2.18	R	9					C	ff
329	Sha00	^{32}S	^{208}Pb	^{240}Cf	180	45.5	20.8	1.31	R	9					C	ff
330	Sha00	^{32}S	^{208}Pb	^{240}Cf	200	63.0	36	1.5	R	9					C	ff
331	Hof94	^{32}S	^{208}Pb	^{240}Cf	200	67				8.8					C	
332	Sha00	^{32}S	^{208}Pb	^{240}Cf	215	76.0	54	1.7	R	9					C	ff
333	Sha00	^{32}S	^{208}Pb	^{240}Cf	230	89.9	61.3	1.84	R	9					C	ff
334	Hof94	^{32}S	^{208}Pb	^{240}Cf	230	93				8.8					C	
335	Sha00	^{32}S	^{208}Pb	^{240}Cf	245	103.0	67	2.0	R	9					C	ff
336	Sha00	^{32}S	^{208}Pb	^{240}Cf	265	120.5	73	2.1	R	9					C	ff
337	Sha00	^{32}S	^{208}Pb	^{240}Cf	285	138.0	78.1	2.27	R	9					C	ff

No.	S_1	E_1	Γ_1	S_2	E_2	Γ_2	FWHM	def.
286	$0.33 f$	11.3 ± 0.2	3.5 ± 0.5	$0.67 f$	14.4 ± 0.7	5.5 ± 0.7		$0.34^{+0.01}_{-0.02}$
287	$0.33 f$	11.4 ± 0.2	5.1 ± 0.5	$0.67 f$	14.8 ± 0.7	8.0 ± 0.7		$0.31^{+0.03}_{-0.02}$
288		14.0 ± 0.2	5.0 ± 0.3					
289		11.7 ± 0.2	3.5 ± 0.3		16.5 ± 0.4	5.5 ± 0.5		0.41 ± 0.04
290	0.33 ± 0.06	11.7 ± 0.3	3.9 ± 0.4	0.67 ± 0.08	15.0 ± 0.5	4.8 ± 0.5		0.29 ± 0.08
291		11.9 ± 0.1	3.8 ± 0.2	$2S_1 f$	15.4 ± 0.2	5.3 ± 0.3		0.31 ± 0.02
292		11.9 ± 0.1	3.8 ± 0.2	$2S_1 f$	15.4 ± 0.2	5.3 ± 0.3		0.31 ± 0.02
293	$0.33 f$	10.5 ± 0.2	7.0 ± 0.5	$0.67 f$	15.0 ± 0.5	9.0 ± 0.8		0.43 ± 0.05
294	$0.33 f$	11.9 ± 0.2	3.8 ± 0.3	$0.67 f$	15.4 ± 0.4	5.3 ± 0.5		0.31 ± 0.04
295	$0.33 f$	$12.0 f$	$3.9 f$	$0.67 f$	$15.4 f$	$5.4 f$		
296		11.9 ± 0.2	5.8 ± 0.3	$2S_1 f$	$15.8^{+0.5}_{-0.2}$	7.5 ± 0.5		$0.34^{+0.04}_{-0.02}$
297	$0.33 f$	11.5 ± 0.3	6.5 ± 0.4	$0.67 f$	15.8 ± 0.5	8.5 ± 0.5		0.38 ± 0.05
298	$0.67 f$	12.7 ± 0.3	8.4 ± 0.5	$0.33 f$	16.8 ± 0.5	7.1 ± 0.5		-0.31 ± 0.04
299		13.8	9					
300		11.9 ± 0.2	5.8 ± 0.3	$2S_1 f$	15.8 ± 0.3	8.5 ± 0.5		0.34 ± 0.03
301	$0.33 f$	$11.2 f$	$4.5 f$	$0.67 f$	$15.0 f$	$5.5 f$		
302	$0.33 f$	$11.2 f$	$4.5 f$	$0.67 f$	$15.0 f$	$5.5 f$		
303		15.0	5.7 ± 0.5					
304		15.0	6.0 ± 0.5					
305		15.0	6.5 ± 0.5					
306		15.0	6.5 ± 0.5					
307		15.0	7.2 ± 0.5					
308		15.0	7.0 ± 0.5					
309		15.0	8.0 ± 0.5					
310		13.2	7.3					
311		13.2	7.0					
312		13.2	7.0					
313	$0.33 f$	11.0	4.2	$0.67 f$	14.0	5.8		0.3
314	$0.67 f$	12.6	5.5	$0.33 f$	12.8	5.7		-0.2
315	$0.33 f$	11.0	5.7	$0.67 f$	14.0	7.3		0.3
316	$0.67 f$	12.4	6.0	$0.33 f$	13.3	6.9		-0.07
317	$0.33 f$	11.0	6.6	$0.67 f$	14.0	8.2		0.3
318	$0.67 f$	12.3	6.5	$0.33 f$	13.3	7.5		-0.08
319	$0.33 f$	10.3 ± 0.3	4.0 ± 0.6	$0.67 f$	14.0 ± 0.6	5.4 ± 1.0		0.37 ± 0.06
320		$11.2 f$	$4.5 f$	$1 - S_1 f$	$12.2 f$	$5.3 f$		-0.10
321	$0.33 f$	11.2 ± 0.3	4.2 ± 0.6	$0.67 f$	14.2 ± 0.6	5.8 ± 1.0		0.30 ± 0.06
322		$11.2 f$	$4.5 f$	$1 - S_1 f$	$12.2 f$	$5.3 f$		-0.10
323	$0.33 f$	11.0	4.2	$0.67 f$	14.0	5.8		0.3
324	$0.67 f$	12.1	5.5	$0.33 f$	13.3	6.5		-0.1
325	$0.33 f$	11.0 ± 0.3	5.0 ± 0.6	$0.67 f$	14.0 ± 0.6	6.5 ± 1.0		0.29 ± 0.06
326		$11.2 f$	$4.5 f$	$1 - S_1 f$	$12.2 f$	$5.3 f$		-0.10
327		$11.2 f$	$4.5 f$	$1 - S_1 f$	$12.2 f$	$5.3 f$		-0.10
328		$11.2 f$	$4.5 f$	$1 - S_1 f$	$12.2 f$	$5.3 f$		-0.10
329		$12.3 f$	$4.5 f$	$1 - S_1 f$	$13.5 f$	$5.3 f$		-0.10
330		$12.3 f$	$4.5 f$	$1 - S_1 f$	$13.5 f$	$5.3 f$		-0.10
331		$12.4 f$	$4.4 f$	$1 - S_1 f$	$13.3 f$	$6.8 f$		-0.07
332		$12.3 f$	$4.5 f$	$1 - S_1 f$	$13.5 f$	$5.3 f$		-0.10
333		$12.3 f$	$4.5 f$	$1 - S_1 f$	$13.5 f$	$5.3 f$		-0.10
334		$12.2 f$	$5.9 f$	$1 - S_1 f$	$13.7 f$	$8.3 f$		-0.14
335		$12.3 f$	$4.5 f$	$1 - S_1 f$	$13.5 f$	$5.3 f$		-0.10
336		$12.3 f$	$4.5 f$	$1 - S_1 f$	$13.5 f$	$5.3 f$		-0.10
337		$12.3 f$	$4.5 f$	$1 - S_1 f$	$13.5 f$	$5.3 f$		-0.10

REFERENCES FOR TABLE

- Ata90 A. Ataç, J. J. Gaardhøje, B. Herskind, Y. Iwata, S. Ogaza, Phys. Lett. B **252**, 545 (1990).
- Bal94 A. Balanda, J. C. S. Bacelar, E. Běták, J. A. Bordewijk, A. Krasznahorka, H. van der Ploeg, R. H. Siemssen, H. W. Wilschut, A. van der Woude, Nucl. Phys. A **575**, 348 (1994).
- Bau98 T. Baumann, E. Ramakrishnan, A. Azhari, J. R. Beene, R. J. Charity, *et al.*, Nucl. Phys. A **635**, 428 (1998); E. Ramakrishnan, T. Baumann, A. Azhari, R.A. Kryger, R. Pfaff, *et al.*, Phys. Rev. Lett. **76**, 2025 (1996).
- Bra05 A. Bracco, O. Wieland AIP Conf. Proc. **802**, 175 (2005).
- Bra89 A. Bracco, J. J. Gaardhøje, A. M. Bruce, J. D. Garrett, B. Herskind, *et al.*, Phys. Rev. Lett. **62**, 2080 (1989).
- Bra95 A. Bracco, F. Camera, M. Mattiuzzi, B. Million, M. Pignanelli, J. J. Gaardhøje, A. Maj, T. Ramsøy, T. Tveter, Z. Żelazny, Phys. Rev. Lett. **74**, 3748 (1995); A. Bracco, F. Camera, Z. Phys. A **349**, 213 (1994).
- Bra96 A. Bracco, F. Camera, M. Mattiuzzi, D. Gnaccolini, B. Million, J. J. Gaardhøje, A. Maj, T. Tveter, Nucl. Phys. A **599**, 83c (1996).
- Bru88 A. M. Bruce, J. J. Gaardhøje, B. Herskind, R. Chapman, J. C. Lisle, F. Khazaie, J. N. Mo, P. J. Twin, Phys. Lett. B **215**, 237 (1988).
- But90 R. Butsch, M. Thoennessen, D. R. Chakrabarty, M. G. Herman, P. Paul, Phys. Rev. C **41**, 1530 (1990).
- Cam03 F. Camera, A. Bracco, V. Nanal, M. P. Carpenter, F. Della Vedova, *et al.*, Phys. Lett. B **560**, 155 (2003).
- Cam99 F. Camera, A. Bracco, S. Leoni, B. Million, M. Mattiuzzi, *et al.*, Phys. Rev. C **60**, 014306 (1999); F. Camera, A. Bracco, G. Colombo, S. Leoni, B. Million, *et al.*, Nucl. Phys. A **649**, 115c (1999).
- Cha87a D. R. Chakrabarty, S. Sen, M. Thoennessen, N. Alamanos, P. Paul, R. Schicker, J. Stachel, J. J. Gaardhøje, Phys. Rev. C **36**, 1886 (1987).
- Cha87b D. R. Chakrabarty, M. Thoennessen, N. Alamanos, P. Paul, S. Sen, Phys. Rev. Lett. **58**, 1092 (1987); Dipak R. Chakrabarty, Nucl. Phys. A **482**, 81c (1988).
- Cha88 D. R. Chakrabarty, M. Thoennessen, S. Sen, P. Paul, R. Butsch, M. G. Herman, Phys. Rev. C **37**, 1437 (1988).
- Cha96 D. R. Chakrabarty, V. M. Datar, R. K. Choudhury, B. K. Nayak, Y. K. Agarwal, C. V. K. Baba, M. K. Sharan, Phys. Rev. C **53**, 2739 (1996).
- Di600a I. Diószegi, N. P. Shaw, A. Bracco, F. Camera, S. Tettoni, M. Mattiuzzi, P. Paul, Phys. Rev. C **63**, 014611 (2000).
- Di600b I. Diószegi, N. P. Shaw, I. Mazumdar, A. Hatzikoutelis, P. Paul, Phys. Rev. C **61**, 024613 (2000).
- Di601 I. Diószegi, I. Mazumdar, N. P. Shaw, P. Paul, Phys. Rev. C **63**, 047601 (2001).

- Di692 I. Diószegi, D. J. Hofman, C. P. Montoya, S. Schadmand, P. Paul, Phys. Rev. C **46**, 627 (1992); R. Butsch, D. J. Hofman, C. P. Montoya, P. Paul, M. Thoennessen, Phys. Rev. C **44**, 1515 (1991).
- Dre95 Z. M. Drebi, K. A. Snover, A. W. Charlop, M. S. Kaplan, D. P. Wells, D. Ye, Y. Alhassid, Phys. Rev. C **52**, 578 (1995).
- End92 G. Enders, F. D. Berg, K. Hagel, W. Kühn, V. Metag, *et al.*, Phys. Rev. Lett. **69**, 249 (1992).
- Fel93 G. Feldman, K. A. Snover, J. A. Behr, C. A. Gossett, J. H. Gundlach, M. Kicińska-Habior, Phys. Rev. C **47**, 1436 (1993).
- Fli91 S. Flibotte, B. Haas, P. Taras, H. R. Andrews, D. C. Radford, D. Ward, Nucl. Phys. A **531**, 205 (1991).
- Fli96 S. Flibotte, M. Cromaz, J. DeGraaf, T. E. Drake, A. Galindo-Uribarri, *et al.*, Phys. Rev. C **53**, R533 (1996).
- Gaa84 J. J. Gaardhøje, C. Ellegaard, B. Herskind, S. G. Steadman, Phys. Rev. Lett. **53**, 148 (1984).
- Gaa86 J. J. Gaardhøje, C. Ellegaard, B. Herskind, R. M. Diamond, M. A. Deleplanque, G. Dines, A. O. Macchiavelli, F. S. Stephens, Phys. Rev. Lett. **56**, 1783 (1986).
- Gaa88 J. J. Gaardhøje, A. M. Bruce, B. Herskind, Nucl. Phys. A **482**, 121c (1988).
- Gar83 E. F. Garman, K. A. Snover, S. H. Chew, S. K. B. Hesmondhalgh, W. N. Catford, P. M. Walker, Phys. Rev. C **28**, 2554 (1983); A. Lazzarini, D. Habs, W. Hennerici, R. Kroth, J. Schirmer, and V. Metag, Phys. Rev. Lett. **53**, 1045 (1984).
- Ge100 N. Gelli, F. Lucarelli, M. Cinausero, E. Fioretto, G. Prete, *et al.*, Eur. Phys. J. A **7**, 361 (2000).
- Gos85 C. A. Gossett, K. A. Snover, J. A. Behr, G. Feldman, J. L. Osborne, Phys. Rev. Lett. **54**, 1486 (1985).
- Gun90 J. H. Gundlach, K. A. Snover, J. A. Behr, C. A. Gossett, M. Kicinska-Habior, K. T. Lesko, Phys. Rev. Lett. **65**, 2523 (1990).
- Haa83 B. Haas, D. C. Radford, F. A. Beck, T. Byrski, C. Gehringer, J. C. Merdinger, A. Nourredine, Y. Schutz, J. P. Vivien, Phys. Lett. **120B**, 79 (1983).
- Hec03 P. Heckman, D. Bazin, J. R. Beene, Y. Blumenfeld, M. J. Chromik, *et al.*, Phys. Lett. B **555**, 43 (2003).
- Hof94 D. J. Hofman, B. B. Back, I. Diószegi, C. P. Montoya, S. Schadmand, R. Varma, P. Paul, Phys. Rev. Lett. **72**, 470 (1994).
- Hof98 G. van 't Hof, J. C. S. Bacelar, I. Diószegi, M. N. Harakeh, W. H. A. Hesselink, N. Kalantar-Nayestanaki, A. Kugler, H. van der Ploeg, A. J. M. Plompen, J. P. S. van Schagen, Nucl. Phys. A **638**, 613 (1998).
- Kel97 M. P. Kelly, J. F. Liang, A. A. Sonzogni, K. A. Snover, J. P. S. van Schagen, J. P. Lestone, Phys. Rev. C **56**, 3201 (1997).
- Kel99 M. P. Kelly, K. A. Snover, J. P. S. van Schagen, M. Kicińska-Habior, Z. Trznadel, Phys. Rev. Lett. **82**, 3404 (1999).

- Kic87 M. Kicińska-Habior, K. A. Snover, C. A. Gossett, J. A. Behr, G. Feldman, H. K. Glatzel, J. H. Gundlach, E. F. Garman, *Phys. Rev. C* **36**, 612 (1987).
- Kic90 M. Kicińska-Habior, K. A. Snover, J. A. Behr, G. Feldman, C. A. Gossett, J. H. Gundlach, *Phys. Rev. C* **41**, 2075 (1990).
- Kic92 M. Kicińska-Habior, K. A. Snover, J. A. Behr, C. A. Gossett, J. H. Gundlach, G. Feldman, *Phys. Rev. C* **45**, 569 (1992).
- Kic93 M. Kicińska-Habior, K. A. Snover, J. A. Behr, C. A. Gossett, Y. Alhassid, N. Whelan, *Phys. Lett. B* **308**, 225 (1993).
- Kmi00 M. Kmiecik, A. Maj, A. Bracco, F. Camera, M. Casanova, S. Leoni, B. Million, B. Herskind, R. A. Bark, W. E. Ormand, *Nucl. Phys. A* **674**, 29 (2000); A. Bracco, F. Camera, S. Leoni, *Nucl. Phys. A* **682**, 449c (2001); A. Bracco, F. Camera, S. Leoni, B. Million, A. Maj, M. Kmiecik, *Nucl. Phys. A* **687**, 237c (2001).
- Kmi01 M. Kmiecik, A. Maj, A. Bracco, F. Camera, B. Million, O. Wieland, *Eur. Phys. J. A* **12**, 5 (2001).
- Kró92 W. Królas, A. Maj, P. Bednarczyk, B. Fornal, W. Męczyński, J. Styczeń, M. Ziębliński, *Z. Phys. A* **344**, 145 (1992).
- Luc96 F. Lucarelli, N. Gelli, P. Blasi, M. Cinausero, E. Fioretto, *et al.*, *Z. Phys. A* **355**, 35 (1996).
- Mat95 M. Mattiuzzi, A. Bracco, F. Camera, B. Million, M. Pignanelli, J. J. Gaardhøje, A. Maj, T. Ramsøy, T. Tveter, Z. Żelazny, *Phys. Lett. B* **364**, 13 (1995).
- Mat97 M. Mattiuzzi, A. Bracco, F. Camera, W. E. Ormand, J. J. Gaardhøje, A. Maj, B. Million, M. Pignanelli, T. Tveter, *Nucl. Phys. A* **612**, 262 (1997).
- Nan99 V. Nanal, B. B. Back, D. J. Hofman, G. Hackman, D. Ackermann, *et al.*, *Nucl. Phys. A* **649**, 153c (1999).
- Noo92 R. F. Noorman, J. C. Bacelar, M. N. Harakeh, W. H. A. Hesselink, H. J. Hofmann, *et al.*, *Phys. Lett. B* **292**, 257 (1992).
- Noo94 R. F. Noorman, J. C. Bacelar, M. N. Harakeh, W. H. A. Hesselink, H. J. Hofmann, *et al.*, *Nucl. Phys. A* **574**, 501 (1994).
- Rat03 S. K. Rath, D. R. Chakrabarty, V. M. Datar, Suresh Kumar, E. T. Mirgule, A. Mitra, V. Nanal, H. H. Oza, *Phys. Rev. C* **67**, 024603 (2003); D. R. Chakrabarty, *Nucl. Phys. A* **687**, 184c (2001).
- Sch93 J. P. S. van Schagen, Y. Alhassid, J. C. Bacelar, B. Bush, M. N. Harakeh, *et al.*, *Phys. Lett. B* **308**, 231 (1993).
- Sch95 J. P. S. van Schagen, Y. Alhassid, J. C. S. Bacelar, B. Bush, M. N. Harakeh, *et al.*, *Phys. Lett. B* **343**, 64 (1995).
- Sch96 S. Schadmand, R. Varma, S. R. Banerjee, B. B. Back, D. J. Hofman, C. P. Montoya, P. Paul, *Nucl. Phys. A* **599**, 89c (1996).
- Sha00 N. P. Shaw, I. Diószegi, I. Mazumdar, A. Buda, C. R. Morton, *et al.*, *Phys. Rev. C* **61**, 044612 (2000).
- Sto89a A. Stolk, A. Balanda, M. N. Harakeh, W. H. A. Hesselink, J. Penninga, H. Rijneveld, *Nucl. Phys. A* **505**, 241 (1989).

- Sto89b A. Stolk, M. N. Harakeh, W. H. A. Hesselink, H. J. Hofmann, R. F. Noorman, J. P. S. van Schagen, Z. Sujkowski, H. Verheul, M. J. A. de Voigt, D. J. P. Witte, *Phys. Rev. C* **40**, R2454 (1989).
- Thi88 P. Thirolf, D. Habs, D. Schwalm, R. D. Fischer, V. Metag, *Nucl. Phys. A* **482**, 93c (1988).
- Tho87 M. Thoennessen, D. R. Chakrabarty, M. G. Herman, R. Butsch, P. Paul, *Phys. Rev. Lett.* **59**, 2860 (1987).
- Tho88 M. Thoennessen, D. R. Chakrabarty, R. Butsch, M. G. Herman, P. Paul, S. Sen, *Phys. Rev. C* **37**, 1762 (1988).
- Tho92 M. Thoennessen, J. R. Beene, F. E. Bertrand, C. Baktash, M. L. Halbert, *et al.*, *Phys. Lett. B* **282**, 288 (1992).
- Tho93 M. Thoennessen, J. R. Beene, F. E. Bertrand, C. Baktash, M. L. Halbert, D. J. Horen, D. G. Sarantites, W. Spang, D. Stracener, *Phys. Rev. Lett.* **70**, 4055 (1993).
- Tho95 M. Thoennessen, E. Ramakrishnan, J. R. Beene, F. E. Bertrand, M. L. Halbert, D. J. Horen, P. E. Mueller, R. L. Varner, *Phys. Rev. C* **51**, 3148 (1995).
- Vie89 G. Viesti, M. Anghinolfi, P. F. Bortignon, P. Corvisiero, L. Djilavian, *et al.*, *Phys. Rev. C* **40**, R1570 (1989).
- Vie96 G. Viesti, M. Lops, B. Fornal, D. Bazzacco, P. F. Bortignon, *et al.*, *Nucl. Phys. A* **604**, 81 (1996); M. Cinausero, D. Bazzacco, P. F. Bortignon, G. De Angelis, D. Fabris, *et al.*, *Nucl. Phys. A* **599**, 111c (1996).
- Vie97 G. Viesti, B. Fornal, M. Cinausero, *Phys. Rev. C* **55**, 1594 (1997).
- Voj89 R. J. Vojtech, R. Butsch, V. M. Datar, M. G. Herman, R. L. McGrath, P. Paul, M. Thoennessen, *Phys. Rev. C* **40**, R2441 (1989).